



Pediatric Ophthalmology: Amblyopia (Lazy Eye), Strabismus, and Pediatric Eye Examinations

Dr. Ali Wasef Alkhatib ^{*1}

1. Consultant Ophthalmologist, Primary Health Care Corporation, PHCC, Qatar.

***Correspondence to:** Dr. Ali Wasef Alkhatib, Senior Scientist, PhD, Ophthalmologist, Moscow.

Copyright

© 2023 **Dr. Ali Wasef Alkhatib**. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: 23 June 2023

Published: 01 July 2023

Abstract

The preset study was conducted to review updates of the literature regarding clinical problems related to pediatric ophthalmology including amblyopia (lazy Eye), strabismus, and pediatric eye examinations. The field of pediatric ophthalmology is responsible for the discovery and treatment of vision abnormalities in children, both of which are extremely important obligations that fall under its purview. This subspecialty of pediatrics focuses on the treatment of ocular disorders in children. Both amblyopia and strabismus are prevalent disorders that, if left untreated, can have long-term negative repercussions on a person's vision. Amblyopia is more common than strabismus. The condition known as amblyopia is defined by a lack of eye movement, whereas strabismus is characterized by eyes that are turned inward. It is only via the delivery of routine eye examinations to juvenile patients that it is possible to detect these diseases as well as other abnormalities of the eye at an early stage. This not only paves the way for fast treatment but also provides the best visual outcomes that are achievable. Eye exams should be a priority for both parents and healthcare providers so that children can reach their full potential in terms of their visual development. This will allow children to attain their full potential as adults. Because of this, the child's eyes will be able to grow and develop to their full potential.

Keywords: *Ophthalmology, pediatric, amblyopia, strabismus, pediatric eye examinations.*

Introduction

Pediatric ophthalmology is a specialization of the medical science known as ophthalmology. Its primary focus is on the diagnosis and treatment of eye conditions that are prevalent in children. Amblyopia, often known as "lazy eye," and, also known as "misalignment of the eyes," are two of the most common eye problems encountered in pediatric ophthalmology. It is imperative that children get routine eye exams. Amblyopia is one of the most common eye conditions seen in pediatric ophthalmology (Dong et al., 2023).

According to the estimates that were published by the World Health Organization in 2006, there are around 1.4 million children who are blind out of a total of approximately 19 million children under the age of 15 who suffer from some form of visual impairment (VI) (Resnikoff et al., 2004; WHO, 2020).

The majority of these children live in less developed countries in Asia and Africa, which together account for two thirds of the global total. India alone is responsible for the accounting of about 200,000 of these items. Blindness is a disorder that lasts a lifetime; hence, even though there aren't very many blind children, the affliction will have a significant impact on their lives throughout their entire lives (Frick and Foster, 2003). Taking into account the average life expectancy of 70 years, (Riley, 2005), the estimated number of 'blind years' that will be experienced by these 200,000 children is second only to cataract in terms of the severity of the condition. As a result of this, juvenile blindness has important ramifications not just on an individual level but also on a societal and economic societies (Kulkarni et al., 2018).

According to the findings of numerous studies, vision is both the most essential sense for the process of learning and the major channel via which sensory information is taken in. In addition, the majority of a child's early learning is performed through the utilization of their sense of sight (Thompson and Merino, 2018). Because their sensory development is delayed, infants who are born with a visual impairment experience considerable challenges in the areas of cognitive, physical development, and social interaction. This is because of the visual impairment's effect on their sensory development (Dulay and Murphy, 2002; Thompson and Merino, 2018). If they do not receive the most helpful rehabilitation during the early formative years of their lives, this can put them at risk for emotional issues, social isolation, and behavioral challenges (Murphy et al., 2008; Anthony, 2014).

Despite the fact that the necessity of early intervention programs for visually impaired young children is widely acknowledged, there is a paucity of information regarding the effects these programs have in the real world for visually impaired young children (Sahin, 2017).

Amblyopia

The loss of visual acuity in one or both eyes that is caused by incorrect binocular interaction during the critical phase of visual development and that cannot be connected to any ocular or visual system defect or to refractive error is the clinical definition of amblyopia (Guler and Alis, 2022). This loss of visual acuity can be caused by amblyopia in either eye. Amblyopia can affect people of any age, from infants as young as six months to individuals as old as 60 years (Noorden and Campos, 202). According to the American Academy of Ophthalmology, amblyopia is defined as an interocular difference of two lines or more in a visual acuity table (without naming any particular table), or a visual acuity that is worse than or equal to 20/30 with the best optical correction (Zhao et al., 2010).

With a frequency that ranges from 3% to 6% in prosperous countries, amblyopia is the major cause of impaired visual acuity in both children and adults. This condition has a prevalence in these nations. Both the economy and society are significantly impacted by this situation in a meaningful way (Gunton, 2013). Amblyopic individuals typically have less employment opportunities open to them, and their overall quality of life tends to be of a poorer standard. This can present itself in a number of ways, including a person having less social contact, suffering from cosmetic agony (if combined with strabismus), having low self-esteem, experiencing visual disorientation, and being afraid of losing eyesight in the other eye (Wong, 2012; Webber, 2018).

The adoption of interocular difference of visual acuity as a definition takes into consideration many of the points that concern the various other definitions of amblyopia. These points include a reduction in visual acuity, a functional imbalance between the eyes, and inadequate binocular information input in primary visual cortex. In addition, the adoption of this definition acknowledges that amblyopia can be caused by a variety of factors. These are only a few more examples among many others (Levi, 2012; Birch, 2013).

Amblyopia, often known as "lazy eye," is a neurodevelopmental disorder that can affect either one or both of a person's eyes and can manifest at any age (Thykjaer et al., 2017). It is possible for a person to have amblyopia only in one eye. During a child's first few years of life, an abnormal development occurs in the visual pathways that travel from the eye to the brain. This condition is the result of this abnormal development (Reeves et al., 2022). Amblyopia can be caused by a variety of eye conditions, the most frequent of which are refractive irregularities including nearsightedness, farsightedness, or astigmatism. Strabismus, in which one eye deviates from its natural alignment, is another common cause. Amblyopia may also result from the interaction of more than one of these risk factors (Elflein et al., 2015; Ali et al., 2023).

The effective treatment of amblyopia absolutely requires that the condition be identified in its early stages. It is recommended by the American Association for Pediatric Ophthalmology and Strabismus (AAPOS) that children receive a comprehensive eye exam from either an optometrist or an ophthalmologist prior to the age of 3 (Birch et al., 2021). This exam should be performed at least once every year. This is in line with the suggestions that were provided by AAPOS (Holmes, 2020).

Amblyopia can be treated in a variety of methods, such as by patching the eye that is better able to see in order to compel the eye that is less capable to develop increased visual acuity, by wearing corrective glasses, or by participating in vision therapy. Amblyopia is a condition that affects the ability to see clearly. According to Stewart et al (2013), it has been shown that providing children with amblyopia with treatment that begins early and is maintained over time produces better visual outcomes. It has been established via research and investigation that this is the case.

There is abundant data suggesting that the initial treatment for amblyopia should involve correcting the patient's refractive error. This procedure improves the quality of the image that is seen on the retina by using corrective lenses to fix any refractive errors (Birch et al., 2021). When only optical correction was utilized, amblyopia was healed in approximately one-third of previously untreated amblyopic newborns in less than thirty weeks (Pediatric Eye Disease Investigator Group, 2006, 2012). If an individual's condition does not improve after undergoing refractive surgery alone to treat their amblyopia, then they will require additional treatments for their problem. There are treatments that primarily target the impairment of monocular visual acuity, and there are also treatments that target the dysfunction and suppression of binocular vision. Both of these types of treatments are available (Birch et al., 2021).

Strabismus

Strabismus, also referred to as squint or crossed eyes, is a condition that manifests itself when the eyes are not aligned in the correct position (Argiles et al., 2023). It is possible to diagnose it as esotropia, which means a deviation inward (Osborne et al., 2023), or exotropia, which means a deviation outward, and its symptoms may emerge intermittently or continually (Moradi et al., 2023). Esotropia and exotropia both signify a deviation inward. If strabismus is not treated in a timely manner, it can result in amblyopia. This is due to the fact that the brain may choose to disregard information received from the eye that is misaligned in order to avoid experiencing double vision. If strabismus is treated, however, it is possible to avoid this outcome (Birch et al., 2021).

In order to correctly identify strabismus, a patient's eyes need to be carefully examined. During this examination, the patient's visual acuity should be tested, the patient's eye motions and alignment should be evaluated, and the patient's ability to see with both eyes should be evaluated (Li et al., 2015). The various treatment choices for strabismus will be determined not only by the severity of the condition but also by the underlying cause of the condition (Reeves et al., 2022). Vision therapy, vision correction surgery, eye patches, and corrective eyewear are some of the options that may be available (Scheiman et al., 2018). The research that was conducted in 2022 by Kulkarni et al. found that early intervention is essential for preventing long-term visual impairment as well as the psychosocial implications that are linked with strabismus that has not been addressed (Kulkarni et al., 2022).

Pediatric Eye Examinations

The process of detecting and treating visual disorders that affect children includes performing routine pediatric eye examinations. This step is an extremely crucial component of the procedure (Chen et al., 2023). The American Academy of Ophthalmology (AAO) recommends that children receive their initial complete eye checkup when they are roughly six months old (AAO, 2023). After this, additional evaluations are to be carried out on the child when they reach the age of three and before to their enrollment in school (Burton et al., 2020; Wang et al., 2022).

These examinations are performed in an effort to detect any eye abnormalities, including strabismus, amblyopia, and refractive problems, all of which have the potential to hinder the development of one's eyesight (Gupta et al., 2019; Srivastava et al., 2022).

During a pediatric eye exam, an expert in eye care will evaluate the child's visual acuity, eye alignment, eye motions, and the overall health of the eyes. The examination may make use of highly specialized methods such as retinoscopy, which assesses a child's refractive error without the child's cooperation being required in any way in the process (Burton et al., 2022). When eye problems are recognized in a timely way and allowed for, which allows for early intervention, early intervention can significantly improve a child's visual outcomes and overall quality of life (Kulkarni et al., 2022). Early intervention can dramatically improve a child's visual results and general quality of life (Kulkarni et al., 2022).

Summary

The identification and treatment of visual defects in children are vitally significant responsibilities that fall under the ambit of pediatric ophthalmology. This field of medicine treats children with eye conditions. Both amblyopia and strabismus are common diseases that, if left untreated, can have long-term detrimental implications on a person's vision. Amblyopia is characterized by a lazy eye, whereas strabismus is characterized by crossed eyes. The early detection of these illnesses as well as other eye abnormalities, which lays the way for prompt treatment and ensures the best possible visual outcomes, is only feasible through the administration of routine eye examinations to juvenile patients. Eye exams should be a priority for both parents and healthcare providers in order for children to realize their maximum potential in terms of their visual development. This will allow for optimal growth and development of the child's eyes.

References

1. Ali SG, Wang X, Li P, Jung Y, Bi L, Kim J, Chen Y, Feng DD, Magnenat Thalmann N, Wang J and Sheng B (2023) A systematic review: Virtual-reality-based techniques for human exercises and health improvement. *Front. Public Health* 11:1143947. doi: 10.3389/fpubh.2023.1143947.
2. Anthony TL (2014). Family support and early intervention services for the youngest children with visual impairments. *J Vis Impair Blind.*, 108:514–9. [Google Scholar]

3. Birch EE (2013). Amblyopia and binocular vision. *Prog Retin Eye Res.*, 33:67-84.
4. Birch, E. E., Kelly, K. R., & Wang, J. (2021). Recent Advances in Screening and Treatment for Amblyopia. *Ophthalmology and therapy*, 10(4), 815–830. <https://doi.org/10.1007/s40123-021-00394-7>.
5. Burton, M.J.; Ramke, J.; Marques, A.P.; Bourne, R.R.A.; Congdon, N.; Jones, I.; Ah Tong, B.A.M.; Arunga, S.; Bachani, D.; Bascaran, C.; et al. The Lancet Global Health Commission on Global Eye Health: Vision beyond 2020. *Lancet Glob. Health* 2021, 9, e489–e551. [CrossRef] [PubMed].
6. Chen, G.L.J.; Yam, J.C.S.; Pang, C.C.P. Special Issue “Pediatric Eye Disease: Screening, Causes and Treatment”. *Children* 2023, 10, 654. <https://doi.org/10.3390/children10040654>.
7. Chen, G.L.J.; Yam, J.C.S.; Pang, C.C.P. Special Issue “Pediatric Eye Disease: Screening, Causes and Treatment”. *Children* 2023, 10, 654. <https://doi.org/10.3390/children10040654>.
8. D Osborne, J Richardson-May, K Bush, M Evans, SK West, K May & J Self (2023). Four cases of acute comitant esotropia associated with diffuse intrinsic pontine glioma in children, *Strabismus*, 31:1, 26-30, DOI: 10.1080/09273972.2022.2143823.
9. Dong J, Zhu Z, Xu H, He M (2023). Myopia Control Effect of Repeated Low-Level Red-Light Therapy in Chinese Children: A Randomized, Double-Blind, Controlled Clinical Trial. *Ophthalmology*. 2023 Feb;130(2):198-204. doi: 10.1016/j.ophtha.2022.08.024. Epub 2022 Aug 29. PMID: 36049646.
10. Dulay MF, Murphy C (2002). Olfactory acuity and cognitive function converge in older adulthood: Support for the common cause hypothesis. *Psychol Aging*, 17:392–404. [PubMed] [Google Scholar]
11. Elflein HM, Fresenius S, Lamparter J, Pitz S, Pfeiffer N, Binder H, et al (2015). The prevalence of amblyopia in Germany: data from the prospective, population-based Gutenberg Health Study. *Deutsches Ärzteblatt Int.*, 112:338. doi: 10.3238/arztebl.2015.0338.
12. Fatemeh Moradi, Ali Mirzajani, Mohammad Reza Akbari, Masoud Khorrami-Nejad, Jamileh Abolghasemi & Babak Masoomian (2023) Binocular contrast sensitivity in patients with intermittent exotropia in relation to angle of strabismus and level of compensation, *Strabismus*, 31:1, 1-8, DOI: 10.1080/09273972.2022.2141272.

13. Frick KD, Foster A (2003). The magnitude and cost of global blindness: An increasing problem that can be alleviated. *Am J Ophthalmol.*, 135:471–6. [PubMed] [Google Scholar]
14. Guler Alis M, Alis A (2022). Features of the Choroidal Structure in Children With Anisometropic Amblyopia. *J Pediatr Ophthalmol Strabismus*, 59 (5):320-325. doi:10.3928/01913913-20220103-01.
15. Gunton KB (2013). Advances in amblyopia: What have we learned from PEDIG trials? *Pediatrics*, 131(3):540-547.
16. Gupta, R.; Agrawal, S.; Srivastava, R.M.; Singh, V.; Katiyar, V (2019). Smartphone photography for screening amblyogenic conditions in children. *Indian J. Ophthalmol.*, 67, 1560–1563. [PubMed].
17. Holmes JM (2020). When to screen for amblyopia. *J AAPOS.*, 24:65–66.
18. <https://aoa.uberflip.com/i/807465-cpg-pediatric-eye-and-vision-examination>, retrieved in 16/6/2023.
19. Kulkarni S, Gilbert C, Zuurmond M, Agashe S, Deshpande M (2018). Blinding retinopathy of prematurity in Western India: Characteristics of children, reasons for late presentation and impact on families. *Indian Pediatr.* 55:665–70. [PubMed] [Google Scholar].
20. Kulkarni, S. K., Veer, A. B., Gaikwad, A., & Deshpande, M. (2022). The impact of early intervention on a child with vision loss in infancy - A parental perspective. *Indian journal of ophthalmology*, 70(9), 3377–3381. https://doi.org/10.4103/ijo.IJO_412_22.
21. Levi DM (2012). Prentice award lecture 2011: Removing the brakes on plasticity in the amblyopic brain. *Optom Vis Sci.*, 89(6):827-38.
22. Li SL, Reynaud A, Hess RF, Wang YZ, Jost RM, Morale SE, et al (2015). Dichoptic movie viewing treats childhood amblyopia. *J Am Assoc Pediatr Ophthalmol Strabismus.* 19:401–5. doi: 10.1016/j.jaapos.2015.08.003.
23. M. Argiles, J. Gispets, N. Lup on et al., Impact of strabismus and binocular dysfunctions in the developmental eye movement test and test of visual perception skills: A multicentric and retrospective study, *Journal of Optometry* (2023), <https://doi.org/10.1016/j.optom.2023.04.002>.
24. Murphy JL, Hatton D, Erickson KA (2008). Exploring the early literacy practices of teachers of infants, toddlers, and preschoolers with visual impairments. *J Vis Impair Blind*, 102:133–46. [Google Scholar]

-
25. Pediatric Eye Disease Investigator Group (2006). Treatment of anisometropic amblyopia in children with refractive correction. *Ophthalmology.*, 113:895–903.
 26. Pediatric Eye Disease Investigator Group (2012). Optical treatment of strabismic and combined strabismic-anisometropic amblyopia. *Ophthalmology.*, 119:150–158.
 27. Reeves A, Ojha K, Meddaugh H, Zambrano RM (2022). Short stature, hearing loss, retinitis pigmentosa, and distinctive facies syndrome: a case report. *Am J Med Genet A.* 2022:62964. doi: 10.1002/ajmg.a.62964
 28. Resnikoff S, Pascolini D, Etya'Ale D, Kocur I, Pararajasegaram R, Pokharel GP, et al (2004). Global data on visual impairment in the year 2002. *Bull World Health Organ.* 82:844–51. [PMC free article] [PubMed] [Google Scholar]
 29. Riley JC (2005). Estimates of regional and global life expectancy, 1800–2001. *Popul Dev Rev.*, 31:537–43. [Google Scholar].
 30. Sahin C (2017). Increasing the effectiveness of early intervention practices for young children with visual impairment. *Int J Learn Dev.*, 7:30–43. [Google Scholar]
 31. Scheiman M, Chase C, Borsting E, Mitchell GL, Kulp MT, Cotter SA, et al (2018). Effect of treatment of symptomatic convergence insufficiency on reading in children: a pilot study. *ClinExpOptom.*, 101:585593. <https://doi.org/10.1111/cxo.12682>.
 32. Srivastava, R.M.; Verma, S.; Gupta, S.; Kaur, A.; Awasthi, S.; Agrawal, S (2022). Reliability of Smart Phone Photographs for School Eye Screening. *Children*, 9, 1519. [CrossRef] [PubMed].
 33. Stewart CE, Wallace MP, Stephens DA, Fielder AR, Moseley MJ (2013). The effect of amblyopia treatment on stereoacuity. *J AAPOS.*,17:166–173.
 34. Suhr Thykjaer A, Lundberg K, Grauslund J (2017). Physical activity in relation to development and progression of myopia-a systematic review. *Acta Ophthalmol.*, 95:651–9. doi: 10.1111/aos.13316.
 35. Thompson SD, Merino S (2018). Visual Impairments in Young Children: Fundamentals of and Strategies for Enhancing Development. *Young Exceptional Children*, 21:157–69. [Google Scholar]
 36. Von Noorden G, Campos E. *Binocular Vision and Ocular Motility*. 6 eds. St. Louis, Missouri: Mosby, Inc; 2002.

-
37. Wang, Y.M.; Lu, S.Y.; Zhang, X.J.; Chen, L.J.; Pang, C.P.; Yam, J.C. Myopia Genetics and Heredity. *Children* 2022, 9, 382. [CrossRef] [PubMed].
 38. Webber AL (2018). The functional impact of amblyopia. *Clin Exp Optom.* 101(4):443-50.
 39. Wong AM (2012). New concepts concerning the neural mechanisms of amblyopia and their clinical implications. *Can J Ophthalmol.*, 47(5):399-409.
 40. World Health Organization (2020). Global Initiative for the Elimination of Avoidable Blindness: Action plan 2006-2011. Available from: http://www.who.int/blindness/Vision2020_report.pdf.
 41. Zhao PF, Zhou YH, Wang NL, Zhang J (2010). Study of the wavefront aberrations in children with amblyopia. *Chin Med J (Engl.)*, 123(11):1431-1435.

