

Research Article

Mitochondrial Inner Optic Atrophy 1 (OPA1) Gene is Necessary for Regulating and Activating Lysosome, Related Orphan Receptor A (RORA) Genes, and APOL1 Gene Involved in Autophagy Cells for Anti-Inflammation Processes, where ROR-Alpha Genes Stored as Lysosomal Security Granules Within Autophagy Cells

Ashraf Marzouk El Tantawi*

***Corresponding Author: Dr. Ashraf Marzouk El Tantawi** Biomedical molecular studies. Cancer and viral infections. Canada, Toronto, Goldwin Av. Egypt, Giza, Abd elmonem R.

Received Date: November 11, 2020

Publication Date: December 01, 2020

Abstract

Tumor necrosis factor-alpha(TNF- α) subunits deficiency or the involvement in will lead to Sick cell disease (SCD) which is marked by a phenotypic variability and inflammation plays a major role in SCD pathophysiology, which linked strongly to RORA1 genes expression and functions, and also linked to TNF- α subunits expressions and activities, where TNF- α subunits are so essential for anti-inflammation processes and linked to ROR- α genes activities and functions and necessary for regulation of bone homeostasis in several chronic immune and inflammatory joints and tissue diseases.

The inhibition of TNF α due to inhibition or variations in ROR-alpha genes lead to significant inflammations improvements involved in SCD pathophysiology, and also will lead to increase in Nuclear factor- κ B pathways (NF κ B) catabolic pathways and any remaining of TNF- α will be involved in the NF- κ B signaling pathway due to inhibition in their mitochondrial activities. The inhibition or deficiency in presence of Thymine in the Related Orphan Receptor-A (RORA) genes can reflect a deficiency in mitochondrial synthetase enzyme (where mitochondrial OPA1 gene depending on ribosomal genes activities), that'll lead to down activities in ROR-alpha genes functions, and reductions in TNF- α , TXA2, and in VEGF-A subunits. RORA genes necessary for lipid metabolism where is controlled by mitochondrial phospholipase enzymes productions, similarly, ROR-aGenes are necessary for the promotion and regulations of hepatic glucose metabolism and are necessary for hepatic activities and sestrin-Leu carrier synthesis and activities during hepatic metabolic activities.

Lysosome and ROR-a genes are having so necessary functions of preventing G-protein aggregates associated with neuropathies, and preventing blood platelets aggregations depending on mitochondrial activities through producing its active inflammatory enzymes for acting on any toxic inflammation or any aggregation for producing the active TXA2 subunits which through feedback will reform VEGF-A alpha subunits where can be stored in or as lysosomal secretory granules.

Lysosomes and ROR-alpha genes are also necessary for regulating cholesterol through Conversions cholesterol to bile acids and protecting the liver from fibrosis and inflammation thus RORA genes have strong roles in anti-inflammatory cycles, indicating that strong links in function between lysosomes and RORA genes.

ROR-alpha gene is so necessary for anti inflammations where ROR alpha active gene depending on mitochondrial anti-inflammatory enzymes (synthase, phospholipase, Cox2 enzymes) and regulated by mitochondrial inner membrane OPA1 genes activities which are strongly repaired by acyl-CoA:l-acyl-sn-glycero-3-phosphocholine O-acyltransferase. Where acyl-CoA:l-acyl-sn-glycero-3-phosphocholine O-acyltransferase enzyme is so necessary for re-activating brain acetylcholine and for mitochondrial inner membrane repair and protect the liver from fibrosis.

Reductions in mitochondrial activities will lead to reductions in ROR-alpha gene activities and consequently will lead to reductions in TNF-alpha and in CYP7A gene expression and functions (which are locally produced in the brain, and liver), and will lead to hepatopathy, encephalopathy, a variant of Syndromic Intellectual Disability. Where, ROR-alpha genes are necessary for CYP7A gene productions and activities which are necessary for the conversions of both cholesterol and 27-hydroxycholesterol into bile acids. RORA genes considered are imp for regulating anti-inflammations procedures and necessary for controlling the conversion of cholesterol and 27-hydroxycholesterol into bile acids and are so essential for liver protections and are so necessary for protection from sickle cells diseases.

ROR-alpha has anti-inflammatory effects and is involved in DNA damage, where there a strong link between APOL1 protein and ROR alpha genes within autophagy during their activities upon DNA damage and mitochondria dysfunctions. Where during autophagy activities which contain APOL1 protein (lipoproteins) will need phospholipase enzyme for activating APOL1 gene involved in autophagy cells which expressed from mitochondrial membrane for activating the autophagic APOL1 gene for expressing ROR-alpha genes from the autophagy cells for fast-acting on inflammations, where that previous fact reveal “and I consider it as” that ROR-alpha genes are stored within autophagy cells as active lysosomal security granules for fast-acting on tumors and on inflammations molecules which are involved with the pathogenesis of various disorders, including cancer, neurodegeneration, and inflammatory diseases.

But in the case of deficiency of mitochondrial activities or case of inhibitions of phospholipase enzyme expressed from mitochondrial membrane will lead to un activating APOL1 genes lead to inhibition in releasing RORalpha gene from autophagy cells and then involved in tumor contents and interstitium fluid as inactive molecules.

Keywords

_Endosomes tissue cells, _Lysosome, Tetraspanins,

_(TXA2) Thromboxane-A2, _vascular endothelial growth factor VEGF-A,

_tumor necrosis

Factor-alpha TNF- α subunits, _mitochondrial enzymes,

_Th17 cells, _NR1D1,

_CYP7A genes,

_Related Orphan Receptor-A (ROR-alpha) genes.

_Nuclear factor (NF- κ B),

_acyl-CoA:l-acyl-sn-glycero-3-phosphocholine O-acyltransferase (mitochondrial inner membrane repair enzyme “MIMRE”).

_Apolipoprotein L1 _ APOL1 gene.

_Optic Atrophy 1 (OPA1) gene.

Introduction

Endosome and lysosome compartments are characterized by their morphological and functional properties, that their healthy compositions and nutritious can be healthy for inner cell components or can be un-useful or toxic to living cells depend on its molecular compositions and their main feedback origin.

Endosomes are typically involved with the filtering and delivering of lipid vesicles including sorted digested amino acids to and from the plasma membrane then to inner cell components. ‘Early’ endosomes at young ages are close to sites of active endocytosis, where they can act as recycling or re-functioning and transport signals genes and vesicles budding compartments molecules to and from the plasma membrane (which are formed from inner cells) to other specific specialized

tissue cells. Thus, endosomes are provided with sorted lipid, protein, and extracellular molecules to be filtered and transmitted for cell metabolism.

Endosomes, have the transport growth factors and signals transmission functions along the axon, provide rapid genes signals propagations and regulations of signaling cues.

Endosomes are a type of extracellular Gene that originally in young ages are active micro alpha subunits where can stimulate PPARs genes from plasma for antigen synthesis and for inner cells nourishing, and are responsible for inner cell nutritious and protections including anti-inflammatory effects and more as receiving molecules to be filtered for inner cells metabolism and then transmitted as active signals transmission, but as age increased as the composition of endosomes from fatty acids and amino acids will be increased and contaminated that will include necessary and unnecessary fatty and amino acids and may include +ve cationic molecules that will delay many of endosomes functions and activities.

Intercellular connections through endosomes and lysosome are through producing exosomes for running specific metabolic processes, and exosome can be considered as signals genes where can be transmitted by endosomes, and are controlled by RORA genes functions and by the lysosomal secretory granules activities, which are involved with the pathogenesis of various disorders, including cancer, neurodegeneration, and inflammatory diseases (1).

The overexpression of the Orphan nuclear receptor (TR3) in human umbilical vein endothelial cells (HUVECs) resulted in VEGF-A-independent synthesis, survival, and induction of several cell cycle genes activities (13).

Normal endosomes must contain the main receptors for TXA2 subunits, VEGF-A, and TNF α expressions and activations directly or indirectly, that started by producing mRNA which will follow pathways of productions through translations process with the help of stimulators and activators from other sources and factors as ribosomal ATPase, mitochondrial membrane functions, and lysosomes.

The delay in genes signals transmissions by endosomes can cause tumor synthesis in a specific tissue and can cause capillaries blockage, that can be due to an increase in cationic positive elements and linkages which can cause idle and inactivities to some active sites in the gene, that will increase the possibility of neutral inactive linkages, and will be the source of distributing variations and inflammations to all other tissue.

As age increased as endosomes and exosomes compositions will contain healthy nutritious components contaminated with Inert idle inactive bonding molecules in a little constituent in their structure, that those idle inactive bonds during heat shock some of those bonds will Broken then will rapidly be formed again in sudden formations fashions that their electrons will be fast rearranged to form new active linkages with new higher active bonding energy value to form an active tetra molecule which is the Tetraspanins.

Moreover, an increase in ROR α genes functions will activate AMPK pathways, Upregulated antioxidative and will activate anti-inflammatory genes, which ameliorated the symptoms of NASH in the methionine and choline-deficient diet mouse model (25, 26), indicating that ROR-alpha gene is so necessary for anti inflammations where ROR alpha active sites activate and regulate Thromboxane-A2 productions and VEGF-A expressions which depending on mitochondrial anti-inflammatory enzymes (synthase, phospholipase, Cox2 enzymes) and regulated by mitochondrial inner membrane OPA1 genes activities which are strongly repaired by acyl-CoA:l-acyl-sn-glycero-3-phosphocholine O-acyltransferase. Where acyl-CoA:l-acyl-sn-glycero-3-phosphocholine O-acyltransferase enzyme is so necessary for re-activating brain acetylcholine and for mitochondrial inner membrane repair and re-activities, and consequently necessary for ROR-alpha genes re-expressions.

OPA1 genes is the main gene in mitochondria, that when stimulated for repair inner and outer mitochondrial membrane will cleavage to produce L-OPA1 & s-OPA1 genes, where s-OPA1 is involved in the outer membrane synthesis upon fusion with MFN2 gene for activating mitochondria. Where decreasing mitochondrial activities will cause a variant of Syndromic Intellectual Disability with Either Autism or Cerebellar Ataxia (21).

ROR-alpha genes are necessary for CYP7A synthesis, which are locally produced in the brain and in the liver, where necessary in the brain to regulate the synthesis of neurosteroids (17), wherein the liver is acting for catalyzing the hydroxycholesterol in the regulations of cholesterol synthesis (18). Where a 7 alpha-hydroxylation is necessary for the conversion of both cholesterol and 27-hydroxycholesterol into bile acids, and the RORA genes are necessary for CYP7A gene productions and activities which are necessary for the conversion of both cholesterol and 27-hydroxycholesterol into bile acids. RORA genes considered are imp for controlling the conversion of cholesterol and 27-hydroxycholesterol into bile acids and are so essential for liver protections and are so necessary for protection from sickle cell diseases.

Also, deficiency in ROR- α gene activities will reflect decreasing in CYP7A gene activities and will lead to decreasing in bile acids and lead to decreasing in liver protections from fibrosis and from inflammation (19). Also, the high decrease in RORA genes in the human will lead to dramatic increases in basal expression of NF- κ B regulated genes (20), due to the high decreasing in TNF α and in VEGF-A subunits productions.

Methods

Tetraspanins characteristically containing 4, 6 or 8 conserved cysteine residues (3), which indicate that the Cys (TGT, TGC) is linked to Thr (ACA, ACG) amino acid and is the main imp amino acid for slowing and control Tetraspanins activities in a distinct way where Cys present at the head of the game then Thr at the tail of the gene will bond together for converting the gene to the circled round inactive or in steady position shape, and that bonds only will be broken by phosphorylation process and by increasing polarities by ATPase for practicing their functions.

The Tetraspanins genes activities can be stopped or slow down by constructing that linkages between Cys and Thr amino acids, or and between Pro and Gly, and between Tyr and Leu amino acids respectively, and then will cause Twisting, bending and rotating to the gene to slow down its activities, but to be activated again those linkages between those amino acids have to be broken down to let and allow the gene to be inactive straight linear forms again and that will be done by phosphorylation process whether by ribosomal ATPase or by G-actin ATPase or through MAPK pathways.

Those necessary amino acids Thr, Tyr, Gly, Gly, Leu and Cys are depending on their pyrimidine functions where their re-synthesis are depending on and controlled by mitochondrial synthetase enzyme which regulates the conversion of purines to a pyrimidine.

The presence of active cysteine with Tyr, Gly, Gly Leu and Thr amino acids in the exosomes & in Tetraspanins are so necessary directly or indirectly for VEGF-A subunits re-synthesis, and for TNF α synthesis, consequently for endothelin-1 synthesis (4), where, Cys, Arg, Tyr, Gly, Gly, Phe, Leu, Ser, Thr are necessary for anti-inflammation cycles through the effect of mitochondrial anti-inflammation enzymes (synthase, phospholipase, Cox2) on inflammations molecules to produce TXA2 subunits, which through feedback will re-produce VEGF-A alpha subunits, and then will resynthesis Endothelin-1 during anti-inflammations cycles and processes, then will re-stimulate PPARs genes activities with MAPK pathways for completing the rest of complete

cycles. Also, we know that Thr with Cys and Ser with Arg amino acids are so necessary for completing phosphorylation processes in several genes during MPAK pathways which are controlled and regulated by ribosomal ATPase activities.

Indosomes are imp for regulate metabolic cycles through their specific amino acid arrangements to regulate and running anabolic processes and regulate transmitted genes as signals across its polar active receptors in controlled limited steps depending on ribosomal activities and on surrounding stimulating active sources and factors. As I reported before that the increase in mitochondrial synthase, phospholipase, and cox2 enzymes will lead to increasing in TXA2 alpha subunits expressions which will lead to an increase in the anti-inflammation process than through feedback will lead to increasing for VEGF-A alpha subunits re-synthesis and, then will stimulate PPARs and MPAK pathways activities. As VEGF-A subunits and TNF α synthesized as VEGF-B will start to produce under controlling of their VEGF -alpha subunits which will lead to decreasing in the Nuclear factor (NF- κ B). catabolic Processes, eg in joints, and in muscles.

The more of TXA2 and VEGF-A alpha subunits productions, the more possibilities of can be restored in cells as granules to be ready for safety and security for acting as anti-inflammatory tools on microbe toxic molecules and on inflammations molecules, where can be stored in inner cells as lysosomes secretory granules. T-cells and many specific tissue cells are restored their TXA2 and VEGF-A alpha subunits as lysosomal secretory granules in their inner components.

T Cells kill their targets by releasing very potent active cyto- high energy proteins released from their stored granules which are characterized as lysosomal secretory organelles. The accurate killing is assured by a vectorial stimulation to secretory lysosomes along microtubules and focused secretion within the immunological synapse (7).

Retinoic acid-related orphan receptor α (ROR α) activities have a major role in various biological metabolic processes including the anti-inflammation processes and involved in autophagy cells, where, retinoic acid-related orphan receptor α (ROR α) controls the inflammatory signaling network (15).

RORA genes involved in lipid metabolism where is controlled by mitochondrial phospholipase enzymes (which is necessary for activating APOL1 gene which involved in autophagy cells synthesis) for producing apolipoproteins, Where ROR-alpha genes are produced by autophagy cells for acting on inflammations, and also ROR alpha genes involved in the regulation of hepatic glucose metabolism, where is necessary for hepatic activities and is necessary for sestrin-Leu carrier synthesis and activities during hepatic metabolic activities, that is necessary for PPARs

genes activation. Also ROR-alpha gene activities depending on Thymine nucleotide resynthesis which is regulated by and depending on mitochondrial synthetase enzyme which regulates the pyrimidine synthesis in vivo. Where, the most necessary amino acids in the ROR-alpha gene are Ser, Arg, Tyr, Gly, Gly, Phe Leu, and Thr amino acids, that are necessary for TXA2 and then for VEGF-A subunits productions and activities, where any deletion within the ROR-alpha Gene leads to an over-expression of inflammatory cytokines (8), which reveals the roles of RORA genes activities in regulations the anti-inflammation processes and cycles.

Apolipoproteins is a lipoprotein expressed by the APOL-I gene that forms a trypanosome complex which acts as an anti-inflammatory tool and has hemoglobin-binding functions that controlled by the presence of phosphodiesterase and mitochondrial phospholipase enzyme functions. APOL1 are involved in autophagic cell death in case of inhibitions of mitochondrial phospholipase enzymes activities, where reveal that APOL1 controlled by mitochondrial phospholipase enzymes. Where overabundance of APOL1 in the availabilities of mitochondrial phospholipase enzyme within a cell results in autophagy.

Where Autophagy is important for the maintenance of cytoplasmic homeostasis. But, the orphan nuclear receptor, RORa, was identified as a direct target gene of p53. Upon DNA damage (27). And, at the same time, RORa controls the inflammatory state of human macrophages (28). So, as APOL1 protein is a lipoprotein involved in macrophages synthesis and control inflammation processes and involved in cell death in the cause of dysfunction of mitochondria, and as ROR-alpha has anti-inflammatory effects and involved in DNA damage, as they're a strong link between APOL1 protein and ROR alpha genes during autophagy activities and upon DNA damage and mitochondria dysfunctions. Where during autophagy activities which contain APOL1 protein (lipoproteins) will need phospholipase enz expressed from mitochondrial membrane for activating the autophagic lipoproteins (APOL1 protein) than for producing ROR-alpha genes for acting on inflammations molecules, which I consider and reveal that ROR alpha genes are stored within autophagy cells as lysosomal security granules which are involved with the pathogenesis of various disorders, including cancer, neurodegeneration, and inflammatory diseases in case of deficiency of mitochondrial activities or in case of inhibitions of phospholipase enzymes which expressed from the mitochondrial membrane, which needed for activating APOL1 gene involved in autophagy cells for producing ROR alpha genes for fast-acting on inflammations molecules.

Lysosomes functions are so essential for degrading G-protein aggregates associated with neuropathies such as Huntington's disease (HD) and are important for regulating cholesterol homeostasis (14), where RORA alpha genes are also necessary for regulating cholesterol too

through conversions cholesterol to bile acids and protecting the liver from fibrosis and from inflammation thus RORA genes have strong roles in anti-inflammatory cycles, indicating that links in function between lysosomes and ROR families.

ROR α upregulated antioxidative and anti-inflammatory genes, which ameliorated the symptoms of NASH in the methionine and choline-deficient (25, 26), indicating that ROR-alpha gene active sites have strong relations with mitochondrial inner membrane OPA1 genes which are repaired by acyl-CoA:1-acyl-sn-glycero-3-phosphocholine O-acyltransferase, where that enzyme is so necessary for reactivating brain acetylcholine and for mitochondrial membrane repair and re-activities, consequently, that transferase enzyme necessary for reactivating ROR-alpha genes re-expressions for ameliorated the symptoms of NASH in the methionine and choline-deficient.

Lysosome and RORA genes are both having the necessary functions of preventing G-protein aggregates associated with neuropathies, and preventing blood platelets aggregations depending on mitochondrial optimal activities through producing its active inflammatory enzymes for acting on any toxic inflammation or aggregated molecules to produce active TXA₂ subunits which can be stored in or as lysosomal secretory granules, and then transmitted with mitochondrial synthase enzymes and phospholipase enzymes as active signals genes or subunits throughout plasma membrane and endosomes to interstitium fluid between cells for acting on aggregated platelets or G-proteins and acting on inflammations for producing Thromboxane-A₂ then throughout feedback will produce VEGF-A subunits for farther the rest of anti-inflammation cycles activities for reactivating endothelin₁ and G-actin filaments, and removing any other toxicity and impurities from blood vessels.

Note, the lysis of any aggregations whether blood platelets or G-proteins aggregation in vivo will need the help of the mitochondrial activities to produce its necessary enzymes for acting on the aggregation as synthase, phospholipase, and Cox2 enzymes, where those enzymes are anti-inflammatory enzymes and are so necessary for acting on any inflammations, where their effects on inflammations molecules will produce TXA₂ subunits which through feedbacks will produce VEGF-A subunits for increasing the anti-inflammation processes and for reactivating endothelin₁ re-synthesis and G-actin filaments re-activities.

So, lysosomes and RORA genes functions are linked together and depending on those anti-inflammatory enzymes from mitochondria for lysis the aggregated G-protein, so are having almost same active receptors that can be activated by mitochondrial active enzymes, which are

stimulated by inflammations molecules or by G-actin isoforms or by the Tetraspanins activities, where Tetraspanins Functions are involved in Regulations of Cellular Signaling (23).

When, a decrease in NR1D1 will knockdown cells resulted in the reductions of expressions of lysosomal-associated membrane protein 1, LAMP1, commensurate (15) (that indicating the direct strong links between NR1D1 and lysosomes), and also will be a result of reductions in VEGF-A subunits productions, then reduction in anti inflammations processes.

The overexpression of the Orphan nuclear receptor (TR3) in human umbilical vein endothelial cells (HUVECs) resulted in VEGF-A-independent synthesis, survival, and induction of several cell cycle genes activities (13). The overexpression of Orphan nuclear receptor (TR3) reflects the increase in mitochondrial activities and increase in its synthase, phospholipase, and Cox2 enzymes that reflect increasing of the actions of those enzymes on inflammations molecules that will induce an increase in TXA2 alpha subunits productions, and therefore increasing in its feedback to produce VEGF-A subunits productions, and then increasing in anti-inflammation cycles.

TxA2 induces platelet degranulation, depending on the mitochondrial activity. The alpha granules or subunits can Lysis platelet or G-protein aggregation, and then will produce TXA2 subunits for re-complete its anti-inflammation pathways activities).

RORa genes involved in regulating the Anti-inflammatory state of human macrophages (10), and protect the liver from inflammations and from fibrosis, and has main roles in activating brain functions against inflammation and toxicity through feeding and activating enkephalin leu pentapeptides or Met pentapeptides and activate many of brain processes to be driven to neuron cells.

The role of TNF- α subunits are so important in anti inflammations through the control of the activation and productions and activities of nuclear factor kappa-B (NF- κ B)(1), where TNF- α roles are controlling NK-kB productions and activities without involving in NK-kB pathways.

Where RORa gene overexpression inhibits proliferation and tumorigenesis of glioma cells lines and GSCs via inhibiting TNF- α to mediate NF- κ B signaling pathway (11), where reactivations of regular expressions of ROR-alpha (RORa) genes functions will stimulate mitochondrial functions which will lead to phospholipase productions for activating lipoprotein for generating GTPase and for activating ROR-alpha genes whether in autophagy or in interstitium fluid, the result of reactivation MAPK pathways and PPARs functions for continuing anti-inflammations procedures

lead to increase TNF- α subunits productions, wherein that previous steps will produce TNF- α subunits with preventing them to don't mediate NF- κ B signaling pathway but only will control TGF- β pathways and activities, (the inhibitions of NF κ B uncontrol catabolic pathways will be through preventing TNF- α to mediate and involving in NF κ B pathways, through increasing TNF- α subunits productions and functions that will control TGF- β activities), were increasing in TNF- α subunits will inhibit platelet aggregations and will inhibit the irregular pathogenic tumors in the Perfect mitochondrial and ribosomal idealism activities.

Reductions in TNF- α subunits productions will reflect a deficiency in mitochondria or and ribosomal activities and will lead to Sick cell disease (SCD) which marked by phenotypic variability and inflammation plays a major role in SCD pathophysiology, which linked to RORA1 genes functions and TNF- α subunits activities, where TNF- α subunits are so essential for anti-inflammation processes, and for the regulation of bone homeostasis in several chronic immune and inflammatory joints and tissues diseases.

The inhibition of TNF α has led to significant inflammations improvements involved in SCD pathophysiology, and at the same time the inhibition or mutations in RORA-alpha (ROR α) genes will lead to inhibitions in TNF α subunits and inhibitions in its controlled anabolic processes, and also will lead to increasing in NF κ B catabolic up-normal pathways with involving the remaining TNF- α in NF- κ B signaling pathway.

Also, any variations or mutations or inhibition in ROR-alpha genes can reflect deficiency and inhibition in mitochondrial synthetase enzyme lead to a deficiency in Thymine nucleotides synthesis (which are considered as the Basic configurations of Tyrosine, Leu, Cytosine and many other essential amino acids which are so necessary for basic compositions and configurations of active sites in ROR-alpha (ROR α) genes, TNF- α subunits, TXA2 subunits, and VEGF-A subunits, and can reflect a severe deficiency in synthetase with phospholipase and synthases enzymes from mitochondrial membrane activities.

ROR α genes have the function of activating AMPK, Moreover, ROR α upregulated antioxidative and anti-inflammatory genes, which ameliorated the symptoms of NASH in the methionine and choline-deficient diet mouse model (25, 26). As I mentioned before that ROR α genes have the function of activating AMPK, and also ROR α genes are activated by mitochondrial functions but can re-stimulate to upregulated antioxidative and anti-inflammatory genes activities.

The enzyme acyl-CoA:l-acyl-sn-glycerol-3-phosphocholine O-acyltransferase is so necessary for mitochondrial inner membrane OPA1 genes repair and in the main time is necessary for brain

acetylcholine reactivities, were indicated to me that acyl-CoA:l-acyl-sn-glycerol-3 phosphocholine O-acyltransferase can stimulate and reactivate the ROR-alpha genes, which can reactivate anti-inflammatory and antioxidants processes through activating PPARs genes functions and MAPK pathways.

ROR γ t and ROR α related orphan nuclear receptor (ROR) are expressed in Th17 cells, where The loss of Th17 cell populations has been linked to chronic inflammation and microbial translocation, and those regulatory Th17 cells depend on and controlled by mitochondrial synthase, phospholipase enzymes to regulate anti-inflammations synthesis which Promotes Th17 Cells functions (16), and autoimmunity.

Where Th17 cells are so necessary for the maintenance of mucosal immunity but only are regulated and controlled by mitochondrial membrane functions through producing its anti-inflammatory enzymes (phospholipase, synthase and COX-2 enzymes), which basically regulated by ribosomal genes and ATPase enzymes.

Also, the ROR α gene is a regulator of Treg genes responsible for suppressing allergic skin inflammation (24) through its roles for regulating and participating in anti-inflammation processes.

Localization of TNF- α genes are on chromosome 6 (6p21.3) (12), where is localized just beside lysosomal secretory granules, that its basic activity depending on the mitochondrial activities, where mitochondrial synthetase enzyme regulate the expression of Thymine nucleotides from purines through mitochondrial synthetase enzymes in vivo , and is regulating most or all metabolic cycles directly in males, and indirectly in females, where pyrimidine nucleotides are the main for 1st DNA strand in controlling 2nd DNA strand in males, but pyrimidine is main in the 2nd DNA strand for controlling 1st DNA strand in females where the most nucleotides in 1st DNA strand in females are Gly, Glu, Ala,, Arg, Lys ,Thr, Gln ,Pro, leu(AGC) and a lesser of Ser nucleotides, where if 1st DNA strand in females contaminated with more pyrimidine nucleotides can lead to males characters or layer can lead to variations of normal genes lead to severe health problems including cancer, so when some mutated active sites 2nd DNA strand dominant on 1st DNA strand in males will cause health problems, and also when some variant active sites in 2nd DNA strand in females will dominated on their corresponding 1s DNA strand will cause problems, means mitochondrial synthase enz is controlled and stimulated by their main ribosomal genes which corresponding to the related Gender .

Results and Conclusions

APOL1 protein is a lipoprotein involved in macrophages synthesis and involved in inflammations processes and also involved in cell death in the cause of dysfunction of mitochondria, where ROR-alpha has anti-inflammatory effects and involved in DNA damage. There a strong links between APOL1 protein and ROR alpha genes during autophagy activities and upon DNA damage and mitochondria dysfunctions. Where during autophagy activities which contain APOL1 protein (lipoproteins) will need phospholipase enzyme expressed from mitochondrial membrane for activating the autophagic lipoproteins (APOL1 protein) for producing ROR-alpha genes which involved within autophagy cells for acting on inflammations molecules, which reveal and I consider that ROR-alpha genes are stored within autophagy cells as lysosomal security granules which are involved with the pathogenesis of various disorders, including cancer, neurodegeneration, and inflammatory diseases in the case of deficiency of mitochondrial activities or in case of inhibitions of phospholipase enzymes, which expressed from the mitochondrial membrane, which needed for activating APOL1 gene (which involved in autophagy cells) for producing ROR-alpha genes from autophagy cells for fast-acting on inflammations molecules.

The anti-inflammatory cycles are conserved with high-quality accuracy in healthy tissues and can start as full cycles or less fast processes for acting on foreign molecules and on inflammations molecules, whether through Th17 Cells or through mitochondrial synthase, COX-2, and phospholipase (which is necessary for activating APOL1 gene (the lipoproteins) in autophagy cells for producing ROR alpha genes from autophagy for acting fast on inflammations molecules and on foreign cells) Enzymes for producing Thromboxane-A2 (TXA2) subunits which through feedback will regenerate VEGF-A subunits productions which will control the endothelin-1 resynthesis and consequently, will reactivate G-actin subunits functions.

Over increasing in sulfur and in +ve cationic units in genes & in plasma membrane will be the result of increasing the idle inactive subunits in genes or in plasma, which will be the result of formation the hard and rigid membrane structure, that will delays signals genes transmission between cells and tissues, and will delay metabolic processes pathways including anti inflammations processes, that can be the result of blood clots in vessels, specifically, blockage in the capillaries that will isolate those cells regions tissue from being connected to other cells tissue then will lead to tumor synthesis.

ROR-alpha genes may be containing Tyr, Ser, Arg, Thr, gln, and Leu, where the optimum active sites nucleotides arrangements in RORA straight active genes will be Ser, Tyr, Gly, Gly, Phe, Leu, Arg which are necessary for feeding and re-activating acetylcholine and enkephalin Leu pentapeptides activities in the brain and will be reactivated by the mitochondrial repair enzyme : acyl-CoA:1-acyl-sn-glycero-3-phosphocholine O-acyltransferase which is so necessary for repair and reactivate mitochondrial inner OPA1 gene which performs most of the mitochondrial activities.

Where decreasing in the RORA alpha gene's activities will reflect decreasing in mitochondrial main OPA1 gene activities, and reflect decreasing in the mitochondrial repair enzymes and consequently will reflect decreasing in APOL1 gene activities in autophagy cells.

The sharp reduction in ROR α Genes functions in some pathogenic cases will reflect decreasing in mitochondrial activities and deficiency in mitochondrial enzymes expression and will lead to reductions in TNF α subunits productions, and increasing in the catabolic Nuclear factor(NF- κ B) pathways, that will lead to osteoarthritis (OA), and lead to decreasing in anti-inflammation processes and decreasing in immune efficiency.

ROR α genes are so necessary for proliferation for regulating sestrin synthesis in the liver through mitochondrial activities, but the main enzyme needed to be expressed for previous steps from mitochondria is the synthetase enzyme which necessary for regulating pyrimidines synthesis from purines for regulating Leu synthesis and activities.

The deficiency in synthetase enzyme and dysfunction in mitochondria will lead to dysfunction in ROR α gene and consequently in sestrin-Leu carrier tool, then dysfunction in the conversion of the cholesterol to bile acids through decreasing in CYP7A gene synthesis, which is locally produced in the brain and in the liver, lead to decreasing in catalyzing the hydroxycholesterol in the regulations of cholesterol synthesis, lead to liver fibrosis, and Atherosclerosis, sickle cells disease and more...

Also, as RORA genes functions control VEGF-A productions and functions, the reduction in RORA alpha genes activities will lead to decreasing in endothelin-1 synthesis and functions lead to increasing in vain blockage and increasing in inflammation molecules in arteries and vain with a decrease in lipid digestions and metabolism that will lead also to precipitation of lipids arteries that can lead to Arteriosclerosis, and cardiovascular disease which is linked to sharp reductions in RORA alpha genes functions and full activities.

As the high cholesterol increase as conversion to bile acids decrease as precipitation in arteries walls will increase, and then atherosclerosis and heart disease will be the result, and liver fibrosis will be formed, and failure in anti-inflammation processes will occur.

Patients with diabetes and atrial fibrillation are lacking the ROR-a genes activities and decrease in CYP7A gene synthesis and activities in conclusion RORa genes regulates pathologic angiogenesis that is necessary for TNF α subunits productions, therefore are necessary for anti-inflammation cycles and processes, RORa genes necessary for activating enkephalin Leu pentapeptides were decreasing in RORA alpha gene's activities will cause variants of Syndromic Intellectual Disability.

Active RORa gene in the presence of mitochondrial optimal activities is so imp for protecting the liver from inflammations and from fibrosis, and also are so imp for feeding and activating enkephalin pentapeptides in the brain for protection from Intellectual Disability.

RORA genes are so necessary for CYP7A gene productions and activities for conversion the cholesterol to bile acids through catalyzing 27-hydroxycholesterol molecules to cholesterol than to bile acids.

Lysosomes functions are so essential for degrading G-protein aggregates were depending on mitochondrial synthase and phospholipase enzymes productivity for acting on aggregation which considered as inflammation (means is responsible for TXA2 subunits productions), and are also important for regulating cholesterol too, therefore lysosomes functions are so connected and linked to RORA genes functions.

Overstimulation to mitochondria activities through increasing intake alcoholic molecules will cause Overstimulation to mitochondria to produce synthetase, COX-2, synthase, and phospholipase enzymes, lead to over digestion to lipid and some genes chains, lead to high catabolic processes in tissue cells, and highly digestion to amino acids in some genes, lead to the mission of the characteristics of those amino acids in their genes where lead to high conversion of purine to pyrimidine, and then purines nucleotides will be lesser or may disappear, then Gly, Cys, Arg, Thr, Ser Leu will be mutated to be tri-coded pyrimidine and will be variable in their biological structure, lead to hardening in arteries and myocardium, and can lead to tumors and cancer. If RORA genes due to any known (or unknown) factors will be mutated will lead to a down in liver activities and protection from fibrosis, down in VEGF-A expression, and will decrease the enkephalin pentapeptides activities, and decrease the genes signals transmission, and will lead to reductions in Tetraspanins activities.

I would like to give imp note that, Any active genes can found in vivo as a straight linear chain (active genetic thread), but can be circled or rounded (cell round shape) in a steady the Lethargic or in a quiet state, where can be circled through covalent linkages between Tyr and Leu, and through linkages between Arg (AGG) at the head of gene and Ser (TCC) at the end part of the gene, where at the time of activating that gene by phosphorylation those bonds between Arg and Ser will be broken through the phosphorylation process lead to the return that gene to its active straight form again.

Where the active gene forms are found in straight linear chains in vivo and depend on the stimulations by phosphorylation to be active. That activity which responsible for changing Circ-RNAs forms to linear active RNA L-RNAs is controlled by Ser&Arg linkages and Thr (ACC) & Trp (TGG) linkages & Thr (ACA) & Cys (TGT) linkages, where it's necessary to break those linkages through phosphorylation to convert circ-RNAs to return to its active L-RNAs forms, but in case of deficiency in phosphorylations, the circ-RNA will not be able to penetrate living cell for feeding ribosome, and will be present in cytosol as tiny spherical organelles, where that decrease of previous mechanisms can lead to delaying in some metabolic processes and can lead to pathogenic diseases eg un-abilities of Autophages to be converted to its L-active forms that will force them to release their active security lysosomal ROR-a gene for acting on inflammation and helping for running other metabolic processes.

Conflict of Interest Statement

The authors declare that the research work was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. "The biology, function, and biomedical applications of exosomes". Science 07 Feb 2020: Vol. 367, Issue 6478, eaau6977 DOI: 10.1126/science.aau6977.
2. "ROR1 sustains multivesicular endosomes by interacting with HRS and STAM1". DOI :10.21203/rs.3.rs-52722/v1.
3. "Structural organization and interactions of transmembrane domains in tetraspanin proteins" Oleg V Kovalenko, Douglas G Metcalf, [...], and Martin E Hemler Published online 2005 Jun 28. Doi: 10.1186/1472-6807-5-11 PMID: 15985154.

4. "Molecular and Cellular Endocrinology" William G. Haynes, David J. Webb, in Principles of Medical Biology, 1997.
5. "Advances in Pharmacology" Volume 77, 2016, Pages 177-208 Chapter Six – Endothelium-Dependent Contractions: Prostacyclin and Endothelin-1, Partners in Crime? Author links open overlay panel O. Baretella P. M. Vanhoutte <https://doi.org/10.1016/bs.apha.2016.04.006>
6. "Protein sorting into multivesicular endosomes" Camilla Raiborg et al. Curr Opin Cell Biol. 2003 Aug. Cell Biol. 2003 Aug;15(4):446-55. Doi: 10.1016/s0955-0674(03)00080-2. PMID: 12892785.
7. "CTL secretory lysosomes: biogenesis and secretion of a harmful organelle". Author : panel Giovanna Bossi Gillian M.Griffiths <https://doi.org/10.1016/j.smim.2004.09.007>.
8. "Increased Ischemia-Induced Angiogenesis in the Staggerer" Mouse, a Mutant of the Nuclear Receptor Ror Sandrine Besnard, Jean-Sébastien Silvestre, Micheline Duriez, Joëlle Bakouche, Yolande Lemaigre-Dubreuil, Jean Mariani, Bernard I. Levy, Alain Tedgui. DOI: 10.1161/hh2401.101755
9. "Rucker D, Dhamoon AS. Physiology, Thromboxane A2. [Updated 2020 Sep 13]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan.
10. "ROR α controls inflammatory state of human macrophages". PLoS ONE 13(11): e0207374. Doi:10.1371/journal.pone.0207374
11. Doi: 10.1016/j.ebiom.2020.102651 PMID: 32062354 Doi: 10.1016/j.ebiom.2020.102651 PMID: 32062354
12. "The Role of TNF-Alpha in ALS: New Hypotheses for Future Therapeutic Approaches" "January 2012 In book: Amyotrophic Lateral Sclerosis Publisher: InTechEditors: Martin H. Maurer.
13. "Orphan nuclear receptor TR3/NR4A1 regulates VEGF-A-induced angiogenesis through its transcriptional activity" April 2006 Journal of Experimental Medicine 203(3):719-29 DOI: 10.1084/jem.20051523 SourcePubMed
14. "Lysosome and Endosome Organization and Transport in Neurons" J.G. Gindhart, K.P. Weber, in Encyclopedia of Neuroscience, 2009.

15. “ROR α is crucial for attenuated inflammatory response to maintain intestinal homeostasis” Se Kyu Oh, Dongha Kim, Kyeongkyu Kim, Kyungjin Boo, Young Suk Yu, Ik Soo Kim, Yoon Jeon, Sun-Kyoung Im, Su-Hyung Lee, Ji Min Lee, Younhee Ko, Ho Lee, Daechan Park, Sungsoon Fang, and Sung Hee Baek PNAS October 15, 2019 116 (42) 21140-21149; first published September 30, 2019; <https://doi.org/10.1073/pnas.1907595116>.
16. “TNF α Promotes Th17 Cell Differentiation through IL-6 and IL-1 β Produced by Monocytes in Rheumatoid Arthritis” Volume 2014 | Article ID 385352 | <https://doi.org/10.1155/2014/385352>.
17. “Cyp7b, a novel brain cytochrome P450, catalyzes the synthesis of neurosteroids 7 α -hydroxy dehydroepiandrosterone and 7 α -hydroxy pregnenolone” Ken A. Rose, Genevieve Stapleton Proc Natl Acad Sci U S A. 1997 May 13; 94(10): 4925–4930. Doi: 10.1073/pnas.94.10.4925 PMID: PMC24607 PMID: 9144166.
18. Martin KO, Reiss AB, Lathe R, Javitt NB. “7 α -hydroxylation of 27-hydroxycholesterol: biologic role in the regulation of cholesterol synthesis”. J Lipid Res. 1997 May;38(5):1053-8. PMID: 9186922.
19. “Cholesterol 7 α -hydroxylase protects the liver from inflammation and fibrosis by maintaining cholesterol homeostasis[S]” Hailiang Liu, Preeti Pathak, [...], and John Y. L. Chiang. J Lipid Res. 2016 Oct; 57(10): 1831–1844. Doi: 10.1194/jlr.M069807 PMID: PMC5036364 PMID: 27534992.
20. “ROR α controls inflammatory state of human macrophages” Neda Nejati Moharrami, Erlend Bjørkøy Tande, Liv Ryan, Terje Espevik, Victor Boyartchuk. PLoS ONE 13(11): e0207374. Doi:10.1371/journal.pone.0207374.
21. The American Journal of Human Genetics Volume 102, Issue 5, 3 May 2018, Pages 744-759 Article “Dual Molecular Effects of Dominant RORA Mutations Cause Two Variants of Syndromic Intellectual Disability with Either Autism or Cerebellar Ataxia” Author links open overlay panel Claire Guissart 137... Sébastien Küry 2438 <https://doi.org/10.1016/j.ajhg.2018.02.021>.
22. “Nuclear receptor ROR α regulates pathologic retinal angiogenesis by modulating SOCS3-dependent inflammation” Ye Sun, Chi-Hsiu Liu, [...], and Jing Published online 2015 Aug 4. Doi: 10.1073/pnas.1504387112 PMID: PMC4547221

23. *Biol.*, 06 April 2017 | <https://doi.org/10.3389/fcell.2017.00034> “Tetraspanins Function as Regulators of Cellular Signaling” Christina M. Termini and Jennifer M. Gillette*
24. “ROR α -expressing T regulatory cells restrain allergic skin inflammation” Nidhi Malhotra, Juan Manuel Leyva-Castillo, [...], and Raif S. Geha. Doi: 10.1126/sciimmunol.aa06923 PMID: PMC5912895 PMID: 29500225.
25. Han YH, et al. “ROR α decreases oxidative stress through the induction of SOD2 and GPx1 expression and thereby protects against nonalcoholic steatohepatitis in mice”. *Antioxid. Redox Signal.* 2014;21:2083–2094. Doi: 10.1089/ars.2013.5655. [PMC free article] [PubMed] [CrossRef] [Google Scholar].
26. “Liver-specific deletion of ROR α aggravates diet-induced nonalcoholic steatohepatitis by inducing mitochondrial dysfunction” Hyeon-Ji Kim, Yong-Hyun Han, [...], and Mi-Ock Lee Published online 2017 Nov 22. Doi: 10.1038/s41598-017-16077-y PMID: PMC5700103 PMID: 29167529.
27. “Damage; cell death; p53; senescence; response”. *Oncogene* (2013) 32, 3789–3797; doi:10.1038/onc.2012.556; published online 3 December 2012.
28. “ROR α controls inflammatory state of human macrophages” Neda Nejati Moharrami, Erlend Bjørkøy Tande, Liv Ryan, Terje Espevik, Victor Boyartchuk. *PLoS ONE* 13(11): e0207374. Doi:10.1371/journal.pone.0207374. Accepted: October 30, 2018; Published: November 28, 2018.

Volume 1 Issue 4 November 2020

©All rights reserved by Dr. Ashraf Marzouk El Tantawi.