



Research Article

Detection of Some Synthetic Cannabinoids (FUB-AMB and AB-FUBINACA) in Blood and Urine using Gas Chromatography-Mass Spectrometry Liquid-Liquid Extraction

Ahmed M.A. Shihata*, A.M. Elessawy¹

1.Forensic Medicine Authority - Ministry of Justice – Egypt

***Corresponding Author: Ahmed M.A. Shihata**, Forensic Medicine Authority - Ministry of Justice – Egypt.

Received Date: February 05, 2021

Publication Date: March 01, 2021

Abstract

In recent years, various types of synthetic cannabinoids have become widely distributed and are causing social and health problems in most of the world. Synthetic cannabinoids are currently the largest group of new psychoactive substances. Those that have been subjected to legal control are replaced by newer controlled and uncontrolled substances. Some of the most recent synthetic cannabinoids that have been distributed on the market between youth are FUB- AMB and AB-FUBINACA. This study quantified blood and urine of two cases of smoking tobacco mix with AMB-FUB 0.06 - 0.03 ng/mL and 1.7 - 2.9 ng/mL AB-FUB urine and blood respectively.

Keywords: FUB- AMB and AB-FUBINACA - Blood and Urine- GC MS.



Introduction

Synthetic cannabinoids are currently the largest, most diverse and fastest-growing group of new psychoactive substances.

Many of the early synthetic cannabinoids that were synthesized for use in the research were named after either the scientist who first synthesized them or the institution or company where they originated. Now many synthetic cannabinoids are assigned names derived from their chemical names.

the natural cannabinoid with the strongest binding affinity to the CB1 receptor, which is linked to the psychoactive effects or "high" of marijuana (1).

These synthetic analogs often have a greater binding affinity and greater potency to the CB1 receptors. There are several synthetic cannabinoid families (e.g. CP-xxx, WIN-xxx, JWH-xxx, UR-xxx, and PB-xx) classified based on the base structure (2).

Synthetic cannabinoids are a class of molecules that bind to cannabinoid receptors in the body (the same receptors to which THC and CBD attach, which are cannabinoids in cannabis plants). They are designer drugs that are commonly sprayed onto plant matter (3).

Most synthetic cannabinoids are agonists of the cannabinoid receptors. They have been designed to be similar to THC (4).

In the late 2000s, synthetic cannabinoids were identified by laboratories in "Spice" or "K2" and related herbal incense products. The compounds found in the "first generation" of synthetic cannabinoid products were primarily the C8 homologs of the nonclassical cannabinoid CP-47,497 and the aminoalkylindole, JWH-018, both of which are agonists of the CB1 and CB2 receptors. The binding affinity of the synthetic cannabinoids to the CB receptors is dependent on the compound. For example, JWH-018 has a binding affinity for the CB1 receptor that is four times greater than that of THC and approximately ten times higher than THC for the CB2 receptor (5,6).

There are five major categories for synthetic cannabinoids: classical cannabinoids, non-classical cannabinoids, hybrid cannabinoids, aminoalkylindoles and eicosanoids. The indazole carboxamide including APINACA, an adamantyl indazole carboxamide, and AB PINACA, an aminocarbonyl indazole carboxamide, is an example of a new group of synthetic cannabinoids (7). Synthetic cannabinoids are metabolized via cytochrome P450 enzymes resulting in phase I hydroxylated metabolites (8). An alkyl side chain, when present, appears likely to undergo hydroxylation at several positions. Compounds



fluorinated at the 5 positions are also susceptible to oxidative defluorination and hydroxylation (9).

Metabolism of AB-PINACA by human liver microsomes suggested hydroxylation occurred primarily on the pentyl chain (10).

Synthetic cannabinoids were originally synthesized to investigate the endocannabinoid system or for their potential therapeutic benefits, but none progressed to clinical use, they are related to chemicals found in the marijuana plant. Because of this similarity, synthetic cannabinoids are sometimes misleadingly called "synthetic marijuana" (or "fake weed"), and they are often marketed as "safe," legal alternatives to that drug. These compounds are either sprayed on dried, shredded plant material so they can be smoked (herbal incense) or sold as liquids to be vaporized and inhaled in electronic cigarettes and other devices (liquid incense).

The greater addictiveness and more severe adverse effects of synthetic cannabinoids in comparison to marijuana are thought to stem from the fact that many of the synthetic cannabinoids are full agonists to the cannabinoids receptors, CB1 and CB2, compared to THC, which is only a partial agonist (11).

Mechanism of action

Synthetic cannabinoids are referred to as substances with structural features that allow binding to one of the known cannabinoid receptors, CB1 or CB2, present in human cells. The association of synthetic cannabinoids with the above chemical receptors leads to an increase in the concentration of another neurotransmitter known as dopamine.

When the concentration of dopamine in the brain rises, it causes a feeling of deformity after the use of these substances, and as is known at high dopamine. The expansion of the blood vessels also leads to increased body temperature and increased sweating. Dopamine is also responsible for awareness, motivation and feeling. This explains the inability of the abusers to perceive the time or to distinguish the night from the day or even the inability to remember what he was doing and what he did and this may continue to eliminate the effect of the drug. The target analyte in the blood is the parent drug. Most synthetic cannabinoids are further metabolized to hydroxyl-synthetic cannabinoids. These hydroxyl metabolites are excreted in urine almost as glucuronides and should be hydrolysis to obtain the free metabolite.



In contrast to the pharmacological properties of synthetic cannabinoid metabolites, their toxicological properties remain less-well characterized, though some work has been done. In the present study, we developed a gas chromatography-mass spectrometry liquid-liquid extraction to determine FUB-AMB and AB-FUBINACA in the blood and urine.

AMB-FUBINACA

AMB-FUBINACA is the methyl ester analog of AB-FUBINACA, where the terminal amide group of the 1-amino-3-methyl-1-oxobutan-2-yl is replaced with a methyl ester group.

AMB-FUBINACA (also known as FUB-AMB and FUB-MMB) N-[[1-[(4-fluorophenyl)methyl]-1H-indazol-3-yl]carbonyl]-L-valine, methyl ester.

AB-FUBINACA

AB-FUBINACA is classified as an indazole. AB-FUBINACA is based on an indazole core structure where the 1- and 3-positions of the indazole ring system are substituted.

AB-FUBINACA also write AB-FUB N-[[1-[(4-fluorophenyl)methyl]-1H-indazole-3-carboxamide




Figure 1. Chemical structure of AMB- FUBINACA




Figure 2. Chemical structure of AB-FUBINACA

Materials and Methods

Chemicals and Reagents

Standards of the synthetic cannabinoid panel were purchased from Cayman Chemical Co. (Ann Arbor, MI, USA) Ammonium sulfate and hydrochloric acid, aqueous ammonia, Ethyl acetate purchased from Algomhoria Co.

Urine preparation

Hydrolyses 2.5mL of urine was added to 0.3mL of hydrochloric acid (20%) and heated at 60-70°C for 60 min.

Extraction

After cooling, aqueous ammonia (25%) was added to reach a pH of 8–9. The samples were extracted with 10mL of ethyl acetate and centrifuged. The organic phase was evaporated by nitrogen stream at 45 °C or below.



Blood preparation

- 1-To 2ml Blood add 5ml of a saturated solution of ammonium sulfate and drops from HCl (0.1M).
- 2- Vortex for 3 min. then Centrifuge for 5 min then taking supernatant.

Extraction

- 3- Add drops of aqueous ammonia (25%) to a pH of 8–9.
- 4-Add 20ml (Ethyl acetate) Shake the mixture for 4 min. using falcon tube.
- 5-Transfer the organic layer into a glass tube and evaporates it by nitrogen stream at 45C.

Instrumental

Gas chromatography (Agilent 6890) coupled to a mass spectrometry detector (Agilent 5973) with column HP 5-MS, (0.25mm×60m×0.25µm) non-polar chromatographic capillary column (Agilent Technologies).

Column:

Agilent HP-5-MS column ; 0.25mm×60m×0.25µm capillary, 60m×250µm×0.250 µm nominal

Mode: Constant flow

Gas type: Helium

Helium pressure: 27.5 Psi

Helium flow: 1.4 ml/min

The separation was performed by applying for the following thermal program

Oven temperature:

Initial temperature: 100°C, hold time: 2 min, ramp 50°C/min to 280 °C, hold time 14.45: Run time 22.6min.

**Inlet:**

Mode: Splitless

Temperature: 250 °C

Pressure: 27.5Psi

Total flow: 1.4 ml/min

Gas saver: Off

MS spect:

Aux temperature: 280 °C

Resolution mode: SCAN

MS source temp: 230°C

MS quad temp: 150°C

Solvent delay: 5min

Retention time: 14.43min. ± 2%

Resolution mode Ions:

The characteristic ions are 109, 253, 324 and 145

The Selected ion group used for qualification analysis for the analyte to match with GC-MS Cayman Spectral library.

Injector:

Injection volume 2 µL



Syringe size: 10 μ L (Auto or Manual injection)

Result and Discussions

FUB- AMB and AB-FUBINACA are synthetic cannabinoids of the class of substituted indazole-3-carboxamides.

First, the blood and urine were examined for the absence of narcotic drugs and psychotropic substances from the groups of amphetamines and their derivatives, benzodiazepine derivatives, cannabinoids, opioids and cocaine.

Urine's the preferred specimen because the collection of urine samples is easy and non-invasive and the concentrations of analytes are often higher when compared with samples of blood or oral fluid. Moreover, the detection time in urine is longer than blood (days or weeks).

The parent compounds may not be detectable in urine, Hydrolysis with acid is required in the procedure of urine sample preparation for FUB- AMB and AB-FUBINACA and other synthetic cannabis compounds because they were found conjugated form excreted.

Blood used for the determination of the concentrations and ratios of parent and its metabolite that could yield useful information relating to acute or chronic use. but Protein blood precipitation is more useful without precipitation.

Blank control of the sample solutions without the target FUB- AMB and AB-FUBINACA of urine and blood not were detected in the solutions used as control blank.

The characteristic fragment ions and the fragmentation pathways of two synthetic cannabis FUB- AMB and AB-FUBINACA were detected in urine and blood parent structure were analyzed carefully. As shown in **Fig.** (3 and 4, 5 and 6).

Two men old 35 and 44 years smoking synthetic cannabis after put in the tobacco.

Results showed a concentration of AMB-FUB urine 0.06 and blood of AMB-FUB 0.03 ng/mL while urine concentration of AB-FUB 1.7 and 2.9 ng/mL AB-FUB blood.




Figure 3. main fragmentation of AB- Fubinaca in Urine (LLE)




Figure 4. main fragmentation of AB- Fubinaca in Blood (LLE)




Figure 5. main fragmentation of FUB- AMB in Urine (LLE)




Figure 6. main fragmentation of FUB- AMB in Blood (LLE)



Conclusion

The presented study of FUB-AMB and AB-FUBINACA are dangerous to health and may lead to fatal intoxication.

References

1. Rapaka, R.S.; Makriyannis, A., eds. (1987). "Structure-Activity Relationships of the Cannabinoids" (PDF). NIDA Research Monograph. 79 – via U.S. Department of Health and Human Services.
2. "K2: Scary Drug or Another Drug Scare?" Newsweek. 2010-03-03. Retrieved 2018-05-04.
3. Macher, R.; Burke, T.W.; Owen, S.S. "Synthetic Marijuana". FBI: Law Enforcement Bulletin. Retrieved 2018-05-04.
4. Canaert, Annelies; Storme, Jolien; Franz, Florian; Auwärter, Volker; Stove, Christophe P. (2016-11-07). "Detection and Activity Profiling of Synthetic Cannabinoids and Their Metabolites with a Newly Developed Bioassay". *Analytical Chemistry*. 88 (23): 11476–11485. doi:10.1021/acs.analchem.6b02600. ISSN 0003-2700. PMID 27779402.
5. Huffman JW, Zengin G, Wu MJ, Lu J, Hynd G, et al. (2005) "Structure-activity relationships for 1-alkyl-3-(1-naphthoyl)indoles at the cannabinoid CB1 and CB2 receptors: steric and electronic effects of naphthoyl substituents. New highly selective CB2 receptor agonists". *Bioorg Med Chem* 13: 89-112.
6. Aung MM, Griffin G, Huffman JW, Wu M, Keel C, et al. (2000) "Influence of the N-1 alkyl chain length of cannabimimetic indoles upon CB1 and CB2 receptor binding". *Drug Alcohol Depend* 60: 133-140.
7. "Details for Synthetic cannabinoids", www.Unodc.Org. Retrieved 2018-05-04.
8. ai S and Fantegrossi WE (2017) "Pharmacological and toxicological effects of synthetic cannabinoids and their metabolites". *Neuropharmacology of New Psychoactive Substances (NPS) The Science Behind the Headlines*:249-262.
9. Wohlfarth A, Castaneto MS, Zhu M, Pang S, Scheidweiler KB, Kronstrand R and Huestis MA (2015) "Pentylindole/Pentylindazole Synthetic Cannabinoids and Their 5-Fluoro Analogs Produce Different Primary Metabolites: Metabolite Profiling for AB-PINACA and 5FAB-PINACA". *AAPS J* 17:660-677.



10. Takayama T, Suzuki M, Todoroki K, Inoue K, Min JZ, Kikura-Hanajiri R, Goda Y and Toyo'oka T (2014) "UPLC/ESI-MS/MS-based determination of metabolism of several new illicit drugs, ADB-FUBINACA, AB-FUBINACA, AB-PINACA, QUPIC, 5F-QUPIC and alpha-PVT, by human liver microsome". *Biomed Chromatogr* 28:831-838

11. Fantegrossi, William E.; Moran, Jeffery H.; Radominska-Pandya, Anna; Prather, Paul L. (2014). "Distinct pharmacology and metabolism of K2 synthetic cannabinoids compared to Δ^9 -THC: Mechanism underlying greater toxicity?". *Life Sciences*. 97 (1): 45–54. doi: jlfs . ISSN 0024-3205. PMC 3945037. PMID 24084047

Volume 2 Issue 2 March 2021

©All rights reserved by Ahmed M.A. Shihata.