

Research Article

Evaluation of Burnout and Job Stress in Care Worker and Comparison between Front-Line and Second-Line in Care Worker During Coronavirus Epidemic.

Niki Sadeghipor, Babak Heidari Aghdam¹, Sahra Kabiri²

1. Babak Heidari Aghdam, Jam general Hospital, Tehran, Iran.
2. Sahra Kabiri, Jam general Hospital, Tehran, Iran

***Corresponding Author: Niki Sadeghipor**, Jam general Hospital, Tehran, Iran.

Received Date: March 29, 2021

Publication Date: April 05, 2021

Abstract

IMPORTANCE Coronavirus disease (COVID-19) is an infectious disease caused by a newly discovered coronavirus.

OBJECTIVE In this study we aimed to the evaluation of burnout and job stress in care workers and comparison between front-line and second-line care workers during the coronavirus epidemic.

The Census method was used to determine the number of participants in the study. In this study, the researchers conducted their research on all people. They gave the questionnaire to all front-line care workers (nurse, assistant nurse, secretary) second-line care workers (Services, security, chefs and hostesses, facilities) of Jam Hospital, which was 537 people, and 342 questionnaires were filled in by the staff.



MAIN OUTCOMES AND MEASURES We focused on symptoms of Job burnout and job stress in Jam Hospital staff. In this method, demographic data such as job, place of work, gender, age, level of education and work shifts were asked and two questionnaires were used. 1- COPSOQ Persian Questionnaire 2- Stress assessment.

RESULTS In the study, 537 health care workers were asked to participate, 342 respondents (63.6%) completed the survey (242[70.7%] (front-line and) 100 [29.3%] (second-line). The occupational data of nonrespondents were similar to those of respondents (Table 1 in the Supplement). Most participants were women (207[60.2%]), were aged 20 to 40 years (260 [76%]), had an educational level postgraduate (199 [58.1, were work night (193 [56.2%]), Of the 342 responding participants, 242 (70.7%) were front-line health care workers directly engaged in diagnosing, treating, or caring for patients with or suspected to have COVID-19, and 100(29.3%) were second-line. A considerable proportion of front-line participants had symptoms of occupational stress($P=0.03$) & Job burnout) $P=0.08$). Job stress and burnout were higher in front-line staff who were in direct contact with patients with COVID 19 than in staff who were not in direct contact with the patient($P=0.02$).

There was no significant relationship between gender ($P=0.5$), education($P=0.3$), job shift ($P=0.06$) and job stress and burnout.

Introduction

Coronavirus disease (COVID-19) is an infectious disease caused by a newly discovered coronavirus (1). In December 2019 The Chinese city of Wuhan reported a novel virus. This virus spread rapidly throughout the world. On 11 March 2020, WHO announced that the outbreak became a global pandemic (2). From the beginning of the pandemic outbreak until to date (February 22, 2021), the following data emerge from the COVID-19 online dashboard of WHO About 110 million people have been infected and about 2,500,000 died from that disease (3). In this critical situation, health care workers on the front line who are strongly involved in the diagnosis, treatment, and care of patients with COVID-19 are at risk of developing psychological distress and other mental health symptoms (4-8) Although the disease is a risk for all people but front-line health care workers are exposed more than others. Such situations threaten not only physical status but also the mental health of front-line workers, for example, the



disease being unknown, lack of medication to treat, lack of adequate free protective equipment, can deeply affect the mental well-being of workers (9-11). Studies conducted during the SARS and H1N1 flu epidemics show that the lack of personal protective equipment is likely to increase the incidence of diseases in health care professionals, who transmit these diseases to their families, which is one of the reasons for the increased stress. Front line care workers experience high levels of anxiety, psychotic, and post-traumatic disorders due to adverse socio-environmental conditions, such as loss of social status and discrimination. In addition to the problems created by the pandemic, public health strategies, such as mandatory isolation, or quarantine in governments' temporary shelters, or the call for people to return to their original places, and social distancing, increase the feeling of loneliness, leading to mental problems that can contribute to suicide (12- 14)

In this study, we aimed to the evaluation of burnout and job stress in care workers and comparison between front-line and second-line care workers during the coronavirus epidemic.

Methods

Study Design

This study followed the Institute for Work & Health (IWH) reporting guideline. Verbal informed consent was provided by all survey participants before their enrollment. Participants were allowed to terminate the survey at any time they desired. The survey was anonymous, and confidentiality of information was assured.

This study is a cross-sectional, hospital-based survey conducted via a region-stratified, 2-stage cluster sampling from Sep 15, 2020, to Dec 10, 2020. During this period, the total confirmed cases of COVID-19 exceeded 29 million in Word. To compare the interregional differences of mental health outcomes among health care workers in Iran, All hospitals in Tehran were involved. We chose Jam Hospital as a sample. Because Tehran was most severely affected. Hospitals equipped with fever clinics or wards for COVID-19 were eligible to participate in this survey.

Participants

The Census method was used to determine the number of participants in the study. In this study, the researchers conducted their research on all people. They gave the questionnaire to all front-line care workers (nurse, assistant nurse, secretary) second-line care workers (Services, security, chefs and hostesses, facilities) of Jam Hospital, which was 537 people, and 342 questionnaires were filled in by the staff.



Outcomes and Covariates

We focused on symptoms of Job burnout and job stress in Jam Hospital staff.

In this method, demographic data such as job, place of work, gender, age, level of education and work shifts were asked and two questionnaires were used.

1- COPSOQ Persian Questionnaire: Kopstuk Questionnaire by Christensen et al. The work environment is in three versions: long, medium and short, and has been translated into different languages, including French, German, Spanish, Swedish, Chinese, etc., and has been used in many authoritative studies. The middle version in 2011 has been translated into Persian by Asalani et al, its validity and reliability have been evaluated and reported. Of the 10 subscales of the Persian version of the Kopstuk questionnaire which include job insecurity (3 items), quantitative work requirements (3 items), requirements (3 items) Cognitive demands of work (4 items), impact on work (3 items), meaning and value of work (3 items), transparency of job responsibilities (4 items), managerial quality (4 items), sense of sociability (3 items) and Job satisfaction (4 items) which was 34 items in total was used to calculate the score. Dimensions The computational formula of the user guide of this scale was used. Each item has five answer options and scores 0, 25, 50, 75 and 100. The score of each subscale will be independent of the others and will be obtained from the mean of the items of the same subscale. Thus, the range of scores in each subscale varies from 0 to 100. The validity and reliability of the average Persian version of the Kopstuk questionnaire have been confirmed in terms of content, face and structure validity. The results of reliability evaluation were also obtained using Cronbach's alpha and internal correlation of 0.7-0.87 and 0.61-0.84, respectively (15).

2- Stress assess: This questionnaire consists of 20 questions. The participant selects one of the following options. Points are added together.

Not at all	Little bit	Some what	Quite a bit	Very much
1	2	3	4	5

Points are added together. What score may indicate:

0 to 35- low stress:

Stress is fairly well managed in your life. It is important to support your body to continue its healthy response.

35 to 70 –moderate stress

Your body’s response to stress may be getting in the way of normal activities, leaving you feeling depleted.



A personalized program may help counteract the effect of stress on your body.

Above 70- high stress

You may be experiencing prolonged stress, and your body's ability to adapt and cope has been compromised. your body systems need support and strategies targeted specifically for you (16).

The different technical titles of respondents refer to the professional titles certificated by the hospital. Participants were asked whether they were directly engaged in clinical activities of diagnosing, treating, or providing nursing care to patients with elevated temperature or patients with confirmed COVID-19. Those who responded yes were defined as frontline workers, and those who answered no were defined as second-line workers.

Statistical Analysis

Data analysis was performed using SPSS statistical software version 26.0 (IBM Corp). The significance level was set at $\alpha = .05$, and all tests were 2-tailed. The original scores of the 4 measurement tools were not normally distributed and so are presented as medians with interquartile ranges (IQRs). The ranked data, which were derived from the counts of each level for symptoms of Job security Quantitative work demands, Emotional work demands, Cognitive work demands, Impact on work, Meaning and value of work, Transparency of job responsibilities, Managerial quality, Sense of socialization, Job satisfaction and job stress, are presented as numbers and percentages.

The nonparametric Mann-Whitney U test and Kruskal-Wallis test were applied to compare the severity of each symptom between 2 or more groups. To determine potential risk factors for symptoms of Job security Quantitative work demands, Emotional work demands, Cognitive work demands, Impact on work, Meaning and value of work, Transparency of the job responsibilities, Managerial quality, Sense of socialization, Job satisfaction and job stress and the associations between risk factors and outcomes are presented as odds ratios(ORs) and 95%CIs, after adjustment for confounders, including sex, age, marital status, educational level, technical title, place of residence, working position (first-line or second-line), and type of section.

Results

Demographic Characteristics

In the study, 537 health care workers were asked to participate, 342 respondents (63.6%) completed the survey (242[70.7%] (,front-line and)100 [29.3%](second-line)).

The occupational data of nonrespondents were similar to those of respondents (**Table 1** in the



Supplement).

Most participants were women (207[60.2%]), were aged 20 to 40 years (260 [76%]), had an educational level postgraduate (199 [58.1, were work night (193 [56.2%]), Of the 342 responding participants, 242 (70.7%) were front-line health care workers directly engaged in diagnosing, treating, or caring for patients with or suspected to have COVID-19, and 100(29.3%) were second-line (**Table 1**).

Table 1- Demographic and occupational characteristics of responders

Characteristic	Total	Occupation							Section				
		nurse	Nurse assistant	secretary	Chef - hostess	security	Servant	Administrative facilities	Special sections	General sections	emergency	OR	Front line
Overall	342	122	22	12	27	2	29	35	82	43	22	114	100
Sex													
Men	135	28	1	1	12	8	29	35	28	0	21	105	84
Women	208	164	28	20	15	0	0	1	49	28	7	9	16
Education level													
≤ Under graduate	143	0	29	21	28	8	29	34	28	0	14	2	98
≥ Post graduate	122	192	0	0	0	0	0	2	49	43	14	112	2
Shift													
Day	142	93	7	21	18	0	8	28	42	36	21	85	27
Night	123	99	22	0	10	8	21	8	36	7	7	29	73
Age													
25-35	137	92	7	11	20	0	7	0	14	7	14	71	27
36-45	123	57	15	10	7	8	8	18	50	29	7	21	19
> 46	82	43	7	0	0	0	14	17	49	7	7	22	54

The severity of Measurements and Associated Factors

A considerable proportion of front-line participants had symptoms of occupational stress($P=0.03$) & Job burnout) $P=0.08$). Job stress and burnout were higher in front-line staff who were in direct contact with patients with COVID 19 than in staff who were not in direct contact with the patient($P=0.02$).



There was no significant relationship between gender (P=0.5), education(P=0.3), job shift (P=0.06) and job stress and burnout.

Employees' job satisfaction dropped sharply during the COVID 19 epidemic (P=0.04), Employees were severely confused in recognizing job responsibilities(P=0.01), They felt insecure at work(P=0.04), Feelings of worthlessness at work(P=0.02) and lack of positive effect of work(P=0.08) were significant in them. During this period, job stress increased in employees(P=0.03). **(Table2)**

Table 2. Severity categories of psychosocial factors, stress and satisfaction with personal protective equipment

Severity category	Occupational								Sex			Working position			Shift			
	nurse	Nurse assistant	secretary	servant	Administrative-facilities	Chef - hostess	security	P value	Men	Women	P value	Frontline	Second line	P value	P value shift	Long	Day	Night
1- Occupational security														0114				
Low	35/9	25	25	27/5	13	9/3	55/5	0/04	13	15	0/4	2816	2613	0/4	15	11/3	33	
Natural	27/4	15/5	0	36	53/3	25	33/4		29	27/6		1412	3619		36	23/3	33	
High	37/3	59/3	70	36	33	55/5	0		57/6	53/3		5712	3111		48	65	33	
2- Quantitative work demands								0/4			0/2			015	0/5			
Low	22/7	37/5	8/25	31/2	10	0	0		87/25	23/2		3915	3418		18/5	24	25	
Natural	21/7	12/5	50	6/2	20	27/5	55/5		20	23/5		1012	2113		24	20	0	
High	55/5	55/5	41/7	62/5	70	62/5	0		62/7	53		5315	4019		57	55	75	



3-Emotional demands of work														0107				
Low	35/825	35/833/2	10	8/3	66/6	0/05	34/6	35	0/6	3313	1013		0/6	36	37	33		
Natural	27/316/6	0	36	20	25	33/3	24/6	23/6		28	2918			25	22	33		
High	37/358/5	55/536	70	66/60			36/6	39/3		5019	4817			38	40	33		
4-Cognitive demands of work														013				
Low	37/359/3	22	50	46/69/3	33/3	0/1	33	30/3	0/3	3212	2915		0/3	25	34	33		
Natural	21/39/3		15/5				33/3	21/3		1416	2815			20	22	33		
High	49/333/3	78	33/3	33/341/5	55/5		33/3	48/3		5014	4311			55	43	33		
5- Impact on work																		
Low	34/666/6	11	33/3	0	25	66/6		35/3	40		3714	3112			30	36	0	
Natural	25	8/3	0	33/3	5/5	33	0	0/008	23	23/3	0/6	1111	1719	014	0/3	25	28	0
High	32/625	89	33/3	73/341/6	33/3			42	37/3		4818	4513			44	45	100	
6- Value of work														0106				
Low	34/3	58/3	33/3	33/3	0	0	0		24/6	33/3		4119	813		30	0	25/6	
Natural	19/6	8/3	11	25	20	25	33/3	0/02	17/3	19/3	0/3	1219	2518		0/7	32	66	12
High	46	50	55/3	33/3	80	58/3	66/6		36/6	41/3		5014	5915			51	33	44
7- Recognition of job responsibilities	/25																	
Low	40	56/2	8/2	48/7	20	12/5	18/7	0/01	31/7	38/2	0/3	3418	2419	0104	0/6	24	33	66
Natural	21	7/75	12/5	25	31/2	50	12/5		14/5	21/7		19175	2416			22	19	0
High	38	31/25	66/7	18/7	75	50	0		53/7	37/5		4513	3519			53/3	47	33



8- Managerial personality																		
Low	21/5	18/7	8/2	12/5	0	12/5	25	0/12	13/7	20	0/3	3418	2419	0103	0/6	14	22	0
Natural	28/2	31/25	33/2	27	15	12/5	25		22/2	29/5		1917	2416			37	16	66
High	49/7	31/25	58/5	60/5	75	75	50		56/7	50/2		4513	3519			51/5	42	33
9- Sense of socialization																		
Low	44/5	59/3	22	15/5	9/5	33/3	33/3	0/1	23/3	42	0/5	4119	2513	014	0/6	22	27	0
Natural	15	9/3	22	33/3	48	15/5	55/5		12/2	59		1514	3916			23	31	0
High	40/5	33	55/5	50	41/5	51/2	11/2		52	38/3		43	3416			55	58/6	0
10- Job satisfaction														013				
Low	38/2	56/2	28/7	25	5	18/7	50	0/04	24	51	0/3	41	2416	013	0/1	29/2	37	0
Natural	24/2	0	37/5	37/5	20	33/25	0		21	26/7		2015	2216			19/2	24/2	0
High	37/5	43/7	66/7	37/5	75	48	50		55/2	40/2		4913	5216			51/5	39	0
stress																		
Low	41	51/2	43/3	75/8	65/2	61/5	56/6		58/5	42/2		4511	6417	016		49/8	39	
Natural	24/5	17/5	15/8	9/1	25/4	24/3	28/4	0/03	20/2	23/5	0/3	1912	2118		0/08	19/6	22/7	15
High	34/5	31/3	40/9	15	9/4	14/1	15		21/75	33/3		3515	1313			30/6	38/3	35

Risk Factors of Mental Health Outcomes

Risk Factors of Mental Health Outcomes analysis showed that Employees who were at the front line of the hospital indirect exposure to patients with covid19 had higher job stress symptoms(p=0.03), Lower



job satisfaction($p=0.04$), Feeling of lower value($p=0.02$) Feeling of job security($p=0.04$) Lack of transparency Job descriptions and responsibilities($p=0.01$).

Discussion

This cross-sectional survey enrolled 324 respondents and revealed a high prevalence of mental health symptoms among health care workers treating patients with COVID-19 in Iran. Overall, 57%, 53%, 50%, 50%, 48.8%, 50.4%, 45.3%, 46.4%, 43% and 49.3% of all participants reported symptoms of, Feelings of lack of job, quantitative demands of work, emotional demands of work, cognitive demands of work, recognition of job responsibilities, the value of work, effectiveness of work, managerial quality, sense of socialization, job satisfaction and job. Participants were divided into 9 groups (front-line and second-line). Most participants were female, were nurses, were night workers, were educated, were aged 20-40 years, and more worked in the front-line. Secretaries, those are working in the frontline and Facility workers who are working in the second-line reported more severe symptoms in all measurements. Our study further indicated that there is no significant relationship between education, gender, and work shift. Working in the front line was an independent risk factor for worse mental health outcomes in all dimensions. Together, our findings present concerns about front-line employees, 49.5% suffered from burnout and 35% from job stress, among which the highest burnout and stress belonged to secretaries, and in second-line employees, 44.8% suffered from burnout and 13.3% suffered from job stress. The highest burnout belonged to the facilities.

In this study, a significant proportion of participants experienced Burnout and stress symptoms, and more than 49.5% reported Burnout and stress.

The psychological response of health care workers to an epidemic of infectious diseases is complicated. Lack of support in the workplace, lack of transparency in job responsibilities have been reported as the most important causes of stress and burnout (1). Also, other factors such as the increase in suspected patients, lack of adequate personal protective equipment, and the possibility of transmitting the disease to the family have aggravated the psychological problems of employees (17, 18).

Of note, 60.2% of all participants were women, and 56.1% were nurses. Our findings further indicate that there was no significant relationship between gender and the rate of burnout and stress. Frontline nurses treating patients with COVID-19 are likely exposed to the highest risk of infection because of their close, frequent contact with patients and working longer hours than usual (19, 20). Moreover, 67.7% of secretaries suffer from burnout and stress despite not being in direct contact with the patient. Also, 44.8% of second-line care workers have suffered from burnout.



It is true that nurses are in direct contact with the patient and the symptoms of burnout and stress are high in them, but the present study shows that secretaries and staff of the second line are at high risk of burnout and stress which is usually ignored.

Limitations

This study has several limitations. First, it was limited in scope. All participants were from Jam hospital in Tehran. Second, not all employees were interested in completing the questionnaire and 60% completed the questionnaire. Third, due to a large number of questionnaire questions, participants may not have completed several questions accurately. Fourth, due to the prolongation of the epidemic period, the psychological symptoms of the employees may have worsened and it is not possible to follow up. Fifth according to our country's situation we have a lot of deficiency in PPE.

Conclusions

In this study, a front-line and second-line care worker in Jam Hospital in Tehran, where patients with covid19 were on the move, it was found that Front-line care workers have a high rate of burnout and job stress. Also, second-line staff who were not in direct contact with a patient with covid19 but travelled to those wards to do services to the patient had symptoms of burnout and job stress. Support for front-line and second-line staff seems necessary.

References

- 1.Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. "Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019". JAMA Netw Open. 2020; 3 (3): e203976.
- 2.Bai Y, Yao L, Wei T, Tian F, Jin D-Y, Chen L, et al. "Presumed asymptomatic carrier transmission of COVID-19". 2020;323(14):1406-7.
- 3.World Health Organization. "Statement on the second meeting of the International Health Regulations (2005).Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV)". Published feb 22.
- 4.Maunders R, Hunter J, Vincent L, Bennett J, Peladeau N, Leszcz M, et al. "The immediate psychological and occupational impact of the 2003 SARS outbreak in a teaching hospital". 2003;168(10):1245-51.



5. Bai Y, Lin C-C, Lin C-Y, Chen J-Y, Chue C-M, Chou PJPS. "Survey of stress reactions among health care workers involved with the SARS outbreak". 2004;55(9):1055-7.
6. Lee AM, Wong JG, McAlonan GM, Cheung V, Cheung C, Sham PC, et al. "Stress and psychological distress among SARS survivors 1 year after the outbreak". 2007;52(4):233-40.
7. Chua SE, Cheung V, Cheung C, McAlonan GM, Wong JW, Cheung EP, et al. "Psychological effects of the SARS outbreak in Hong Kong on high-risk health care workers". 2004;49(6):391-3.
8. Khalatbari-Soltani S, Cumming RC, Delpierre C, Kelly-Irving MJJECH. "Importance of collecting data on socioeconomic determinants from the early stage of the COVID-19 outbreak onwards". 2020;74(8):620-3.
9. Kang L, Li Y, Hu S, Chen M, Yang C, Yang BX, et al. "The mental health of medical workers in Wuhan, China dealing with the 2019 novel coronavirus". 2020.
10. Zarocostas JJTl. "How to fight an infodemic". 2020;395(10225):676.
11. Sheraton M, Deo N, Dutt T, Surani S, Hall-Flavin D, Kashyap RJPr. "Psychological effects of the COVID 19 pandemic on healthcare workers globally: A systematic review". 2020;292:113360.
12. Giorgi G, Lecca LI, Alessio F, Finstad GL, Bondanini G, Lulli LG, et al. "COVID-19-related mental health effects in the workplace: a narrative review". 2020;17(21):7857.
13. Despoina P, Chrysoula DJIJoN. "Investigation of nurses' mental status during Covid-19 outbreak—a systematic review". 2020;7(1):69-77.
14. Taghilou H, Jafarzadeh Gharajag ZJQJoNM. "Investigating the Relationship between Burnout and Job Performance in the Corona Epidemic from the Perspective of Nurses". 2020;9(4):27-33.
15. Pournik O, Ghalichi L, TehraniYazdi A, Tabatabaee SM, Ghaffari M, Vingard EJMjotIRoI. "Measuring psychosocial exposures: validation of the Persian of the copenhagen psychosocial questionnaire (COPSOQ)". 2015;29:221.
16. Haldeman S, Chapman-Smith D, Petersen DM. "Guidelines for chiropractic quality assurance and practice parameters: proceedings of the Mercy Center Consensus Conference: Jones & Bartlett Learning"; 2004.



17.Chan-Yeung MJJoo, health e. “Severe acute respiratory syndrome (SARS) and healthcare workers”. 2004;10(4):421-7.

18.Sinclair RR, Allen T, Barber L, Bergman M, Britt T, Butler A, et al. “Occupational health science in the time of COVID-19: Now more than ever”. Springer; 2020.

19.Li L, Cheng S, Gu JJJ. “SARS infection among health care workers in Beijing, China”. 2003;290(20):2662-3.

20.Shih F-J, Gau M-L, Kao C-C, Yang C-Y, Lin Y-S, Liao Y-C, et al. “Dying and caring on the edge: Taiwan's surviving nurses' reflections on taking care of patients with severe acute respiratory syndrome”. 2007;20(4):171-80.

Volume 2 Issue 4 April 2021

©All rights reserved by Niki Sadeghipor