



Research Article

Journal of MAR Gastroenterology (Volume 1 Issue 3)

Endoscopic Findings in Patients Presenting with Upper GI Bleeding.

Jawad khan ¹, Saira Nasr Malik* ².

1,2. Police and Services Hospital Peshawar. Khyber Teaching Hospital, Peshawar.

Corresponding Author: Saira Nasr Malik, Police and Services Hospital Peshawar. Khyber Teaching Hospital, Peshawar.

Copy Right: © 2022 Saira Nasr Malik, This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received Date: May 22, 2022

Published Date: June 06, 2022

Abstract

Background and aims;

Upper GI bleeding is one of the commonest emergency in gastroenterology having high morbidity and mortality. It has multiple causes from mild to severe outcome. The aim of this study was to evaluate endoscopic finding in patients presenting with upper GI bleeding.

Methods

This is retrospective study which was conducting in police and services hospital and Khyber teaching hospital Peshawar from February 2021 to november 2021. All the patients presenting with upper GI bleeding was evaluated and reviewed along with their records. Data were analysed and entered using SPSS 20.

Results

Total number of patients were 430(225 males and 205 females) who underwent endoscopy presenting with upper GI bleeding. Main age of study population was 52 years with standard deviation of 15 years. Age of the youngest was 15 years while that of oldest was 85 years. More than half of the patients 224 (52%) were in between 41 to 60 years. The commonest endoscopic finding was esophageal varices 224 (52%), followed by portal hypertensive gastropathy 64(15%), erosive gastritis 43(10%) and fundal varices 34(8%), and. Esophageal varices, gastropathy, gastric ulcers and upper GI malignancies were more in male patients as compared to female patients ($p = 0.039$) presenting with upper GI bleeding. While portal hypertensive Gastropathy, erosive esiphagitis and gastric ulcer were more present in female patients. Variceal bleed has having greater dominancy in middle age group ($p = 0.000$). Whereas erosive gastritis, gastric ulcer, and gastric erosions were more present in increased frequency in older (>65 years) UGIB patients ($p = 0.000$).

Conclusion

UGIB was in higher frequency in male patients as compared to females. In our study, esophageal variceal was the major cause of UGIB in this southasian countries while duodenal ulcer bleed was not as common as it is present in other parts of the world. Variceal bleeding was significantly associated with male patients having middle age groups While bleeding from duodenal ulcer was highly associated with older ages.

Introduction

Upper gastrointestinal bleeding (UGIB) is one of the commonest gastrointestinal emergency presenting in ER having high morbidity and mortality, and associated with high burden of health care resources¹. Population based statistical data is important to get insight into the actual burden on health care resources. Upper GI bleed has many causes in different parts of the world. Epidemiological data are helpful in knowing the burden of the problem, the etiology, morbidity and mortality associated with it. Recent advances in medical treatment has changed the dynamics, etiology and treatment of UGIB. There are very few studies which have been done in Pakistan regarding etiology, morbidity and mortality of GI bleed. In Western countries, bleeding from peptic ulcer disease is still the predominant cause of UGIB^{1,4}. recent studies showing after advancement in medical research mortality and rebleeding rates are going down^{1,2}. However, some data have denied these facts³. These data shows that both mortality and morbidity remained unchanged due to upper GI bleed⁴.

There are some other causes of UGIB as well but in very few patients. These include angiodysplasia, aorto-enteric fistula, Dieulafoy's lesions, and hemobilia etc. Vascular bleed is a major culprit of UGIB associated with high morbidity and mortality. Esophageal varices are usually present in lower esophagus. Varices are usually found in portal hypertension as abnormal dilatation of submucosal veins in cirrhosis-related complication of portal hypertension⁵. In countries like Asia and specially South Asia, esophageal variceal bleed is the commonest and deadly cause of UGIB^{6,7}. In contrast to other parts of the world like West, where peptic ulcer disease is number one cause of upper GI bleed. In our study, we have analyzed data of patients with acute UGIB retrospectively noted the cause, clinical presentation, morbidity and mortality related with GI bleed.

Materials and Methods

This was a retrospective study. Data of 430 patients who were admitted to Police and Services Hospital and Khyber Teaching Hospital from February 2021 to November 2021 with UGIB. The study was approved by the respective Ethical Committee. Patients having age of >14 years were included. Hematemesis and melena were the main symptoms of upper GI bleeding. Data was analyzed based on baseline clinical information, laboratory tests, blood unit transfusions, endoscopy records and findings.

Demographics data (age, gender) and endoscopic findings were noted and analyzed. Data were then entered into SPSS. Statistical data analysis was performed with chi-square. Statistical significance was determined at $p < 0.05$. Analysis of information was done according to age and endoscopic diagnosis and presented in table forms.

Results

In our study, 430 patients presenting with UGIB, in those 225 (52.3%) were males and 205 (47.7%) were females. Male to female ratio was 1.09:1. Mean age was 52 years having standard deviation of 15 years. Males mean age was 49 years while females mean age was 50 years. Age of the youngest patient presented was 15 years while 85 years was oldest patient was 85 year. Majority of patients belonged to the fifth decade. About half of the patients 224(52%) having UGIB were having age between 41-60 years.

| Characteristics | N% |
|-------------------------|------------|
| Age(years) | |
| mean | 52 |
| Standard deviation | 15 |
| Range | 14-85 |
| Gender | |
| male | 225(52.3%) |
| female | 205(47.7%) |
| Age group(years) | |
| <40 | 100(23.2%) |
| 41-60 | 224(52%) |
| >60 | 106(24.7%) |

Demographic features in patients presenting with upper GI bleeding.

| Endoscopic findings | Total% |
|----------------------------|---------------|
| Esophageal varices | 224(52%) |
| gastropathy | 64(15%) |
| gastritis | 43(10%) |
| Fundal varices | 34(8%) |
| Duodenal ulcer | 17(4%) |
| Esophageal ulcer | 15(3.4%) |
| Gastric ulcer | 10(2.3%) |
| Gastric erosion | 8 (1.86%) |
| duadenitis | 6(1.39%) |
| Gastric tumor | 4(0.93%) |
| Esophageal erosions | 2(0.46%) |

| | |
|--------------------|----------|
| esophagitis | 1(0.23%) |
| Dieulefoy,s lesion | 1(0.23%) |
| Esophageal tumor | 1(0.23%) |

Endoscopic findings distribution is patients presenting with upper GI bleeding.

Table 2 shows the endoscopic findings in patients of UGIB in our study. esophageal varices was the predominant findings on endoscopy occurring in more than 50% of the patients admitted with UGIB 52% (n = 224). portal hypertensive gastropathy was on 2nd number after esophageal variceal bleed diagnosed in 64 (15%) patients. There were some other minor causes of upper GI bleed, in decreasing order of frequency by erosive gastritis (43, 10%), fundal varices (34, 8%), duodenal ulcer (17, 4%) and other very rare causes (Table 2). If we compare distribution of endoscopic findings according to gender, esophageal varices, erosive gastritis, gastric ulcers and upper GI malignancy were more present in male patients as compared to female patients presenting with upper GI bleed. Portal hypertensive Gastropathy, erosive esophagitis and peptic ulcer were more likely to be present in female UGIB patients.

Discussion

In our study, mean age of presentation was of 52 years which is on lower side as compared to western countries mean age of UGIB patients 7,8. This also shows high rate of aged population in western parts. In our study, UGIB was more commonly diagnosed in male patients when compared to females, having a male to female ratio of 1.09:1. This was comparable to the data reported in western studies in which male patient ratio was greater than females^{2,9}. The commonest cause of UGIB found in this study was upper GI bleeding secondary to esophageal varices (52%). It was followed by other causes like portal hypertensive gastropathy, gastritis, gastric erosions and peptic ulcer. Peptic ulcer is more common in west as compared to east than esophageal varices. the diagnosis of peptic ulcer (gastric and duodenal ulcer) as cause of GI bleed were found in 6.3% of patients in our study which is very low as compared to west. Other local and international studies shows the exact results as well^{10,12}. In contrast to our study findings, several other studies showed peptic ulcer disease as the commonest culprit of UGIB ^{13,16}. High number of esophageal varices in south asian people like India and Pakistan is mainly due to the high prevalence of viral hepatitis (Hepatitis B and C) related cirrhosis. Comparable results to our study are also found in other local studies as well. In a local study, 53% of cases had esophageal varices and 20% had peptic ulcer disease¹⁷. In our study, portal hypertensive gastropathy was the second most common cause of UGIB while in another local study, 64% of cases had esophageal varices and gastric erosions were the second most common cause, i.e., erosive gastritis 15% and PUD 10% of cases ¹⁸.

There were some limitations in our study. As our study was carried out at a tertiary care centre, it may not be representative of the general population. Many patients with milder symptoms may not report to the hospital. Another important thing is that as our hospital is a government run setup, patients mostly belonging to low socioeconomic status come here. Multicenter studies targeting clinics providing EGD services should also be taken into account in future studies to further improve our understanding of different etiologies of UGIB.

Conclusions

It is very essential to have a basic understanding of different conditions causing UGIB. Our study showed that patients presenting with UGIB was more prevalent in male patients. esophageal variceal bleed was the major significant cause of UGIB and peptic ulcer bleed was not as common when compared to the western parts . Esophageal varices, erosive gastritis, peptic ulcers and upper GI malignancies were predominantly found in male UGIB patients. Conversely, female patients presenting with UGIB, portal hypertensive gastropathy, erosive esophagitis and gastric ulcer were major findings on EGD. Esophageal variceal bleeding was mainly a diagnosis in middle age patients. While erosive gastritis, peptic ulcer, gastric erosions and peptic duodenitis were more present in older patients as compared to older patients.

References

1. Loperfido S, Baldo V, Piovesana E, Bellina L, Rossi K, Groppo M, et al. Changing trends in acute upper GI bleeding: A population-based study. *Gastrointest Endosc* 2009; 70:212-24.
2. Di Fiore F, Leclaire S, Merle V, Hervé S, Duhamel C, Dupas JL, et al. Changes in characteristics and outcome of acute upper gastrointestinal haemorrhage: A comparison of epidemiology and practices between 1996 and 2000 in a multicentre French study. *Eur J Gastroenterol Hepatol* 2005; 17:641-7.
3. Thomopoulos KC, Vagenas KA, Vagianos CE, Margaritis VG, Blikas AP, Katsakoulis EC, et al. Changes in aetiology and clinical outcome of acute upper gastrointestinal bleeding during the last 15 years. *Eur J Gastroenterol Hepatol* 2004; 16:177-82.
4. Rockall TA, Logan RF, Devlin HB, Northfield TC. Incidence of and mortality from acute upper gastrointestinal haemorrhage in the United Kingdom. Steering Committee and members of the National Audit of Acute Upper Gastrointestinal Haemorrhage. *BMJ* 1995; 311:222-6.
5. Laine L, Jensen DM. Management of patients with ulcer bleeding. *Am J Surg*. Ahmed A, Ali L, Shehbaz L, Nasir S, Rizvi SRH, Aman MZ, Ali Z: The prevalence of acute upper gastrointestinal bleeding and the

factors causing hemorrhage as observed at a tertiary health care centre in Karachi, Pakistan. *Pak J Surg.* 2017, 33:36-40.

6. Hadayat R, Jehangiri AU, Gul R, Khan AN, Said K, Gandapur A: Endoscopic findings of upper gastrointestinal bleeding in patients with liver cirrhosis. *J Ayub Med Coll Abbottabad.* 2015, 27:391-394.

7. Theocharis GJ, Thomopoulos KC, Sakellaropoulos G, Katsakoulis E, Nikolopoulou V: Changing trends in the epidemiology and clinical outcome of acute upper gastrointestinal bleeding in a defined geographical area in Greece. *J Clin Gastroenterol.* 2008, 42:128-133. 10.1097/01.mcg.0000248004.73075.ad

8. Zaltman C, Souza HS, Castro ME, Sobral M de F, Dias PC, Lemos V Jr: Upper gastrointestinal bleeding in a Brazilian hospital: a retrospective study of endoscopic records. *Arq Gastroenterol.* 2002, 39:74-80. 10.1590/S0004-28032002000200002

9. Mustapha SK, Ajayi N, Jibrin YB, Shehu A: Aetiology of upper gastrointestinal bleeding in North-Eastern Nigeria: a retrospective review of endoscopic findings. *Nigerian J Gastroenterol Hepatol.* 2009, 1:75-78.

10. Dewan KR, Patowary BS, Bhattarai S: A study of clinical and endoscopic profile of acute upper, gastrointestinal bleeding. *Kathmandu Univ Med J.* 2014, 12:21-25. 10.3126/kumj.v12i1.13628

11. Suba M, Ayana SM, Mtabho CM, Kibiki GS: The aetiology, management and clinical outcome of upper gastrointestinal bleeding among patients admitted at the Kilimanjaro Christian Medical Centre in Moshi, Tanzania. *Tanzan J Health Res.* 2010,

12. 12:302-305. 10.4314/thrb.v12i4.51073 12. Sarwar S, Dilshad A, Khan AA, Alam A, Butt AK, Tariq S, Ahmad I: Predictive value of Rockall score for rebleeding and mortality in patients with variceal bleeding. *J Coll Physicians Surg Pak.* 2007, 17:253-256. 05.2007/JCPSP.253256

13. Botianu A, Matei D, Tantau M, Acalovschi M: Mortality and need of surgical treatment in acute upper gastrointestinal bleeding: a one year study in a tertiary center with a 24 hours / day 7 days / week endoscopy call. Has anything changed?. *Chirurgia (Bucur).* 2013, 108:312- 318.

14. Singh SP, Panigrahi MK: Spectrum of upper gastrointestinal hemorrhage in coastal Odisha . *Trop Gastroenterol.* 2013, 34:14-17.

15. Gurung RB, Joshi G, Gautam N, Pant P, Pokhrel B, Koju R, Bedi TR: Upper gastro intestinal bleeding: aetiology and demographic profile based on endoscopic examination at Dhulikhel hospital, Kathmandu University Hospital. *Kathmandu Univ Med J.* 2010, 8:208-211. 10.3126/kumj.v8i2.3560

16. Gaudong Mbethe GL, Mounguengui D, Ondounda M, et al.: Epidemiology of upper gastrointestinal bleeding in Gabon. (Article in French). *Med Sante Trop.* 2014, 24:441-443. 10.1684/mst.2014.0376
17. Pasha MB, Hashir MM, Pasha AK, Pasha MB, Raza AA, Fatima M: Frequency of esophageal varices patients with upper gastrointestinal bleeding. *Pak J Med Sci.* 2011, 27:277-281.
18. Shah MA, Butt Z, Younis I, Afzal M, Atta H, Nadir A: Etiology of upper gastrointestinal bleed at Aziz Bhatti Shaheed Teaching Hospital Gujarat. *Ann Pak Inst Med Sci.* 2016, 12:80-84.